

Ball polytopes and the Vázsonyi problem

Yaakov S. Kupitz¹, Horst Martini, Micha A. Perles

Abstract

Let V be a finite set of points in Euclidean d -space ($d \geq 2$). The intersection of all unit balls $B(v, 1)$ centered at v , where v ranges over V , henceforth denoted by $\mathcal{B}(V)$ is the *ball polytope* associated with V . Note that $\mathcal{B}(V)$ is non-empty iff the circumradius of V is ≤ 1 . After some preparatory discussion on spherical convexity and spindle convexity, the paper focuses on two central themes².

- a) Define the boundary complex of $\mathcal{B}(V)$ (assuming it is non-empty, of course), i.e., define its vertices, edges and facets in dimension 3 (in dimension 2 this complex is just a circuit), and investigate its basic properties³.
- b) Apply results of this investigation to characterize finite sets of diameter 1 in (Euclidean) 3-space for which the diameter is attained a maximal number of times as a segment (of length 1) with both endpoints in V . A basic result for such a characterization goes back to Grünbaum, Heppes and Straszewicz, who proved independently of each other, in the late 1950's by means of ball polytopes, that the diameter of V is attained at most $2|V| - 2$ times, thus affirming a conjecture of Vázsonyi from circa 1935. Call V *extremal* if its diameter is attained this maximal number ($2|V| - 2$) of times. We extend the aforementioned basic result by showing that V is extremal iff V coincides with the set of vertices of its ball polytope $\mathcal{B}(V)$ and show that in this case the boundary complex of $\mathcal{B}(V)$ is self-dual in some strong sense. The problem of constructing new types of extremal configurations will be addressed in a subsequent paper, but we do present here some such new types.

¹Partially supported by the Landau Center at the Mathematics Institute of the Hebrew University of Jerusalem Minerva Foundation, Germany, and by Deutsche Forschungsgemeinschaft.

²For the sake of priority it should be noticed that, in the present form (except for a few chances in the footnotes and one additional reference), this paper was submitted to the journal "Discrete and Computational Geometry" in February 1, 2008.

³This face structure turns out to be quite intricate or, quoting [2], p. 202, lines 22-23: "the face structure of these objects [ball polytopes] is not obvious at all".


MSC Numbers (2000): 52C10, 52A30, 52B10, 52B05, 05C62, 52A40

Keywords: ball hull, ball polytope, barycentric subdivision, canonical self-duality, diameter graph, face complex, face structure, generalized convexity notion, geometric graph, involutory self-duality, Reuleaux polytope, spherical convexity, spindle convexity, Vázsonyi problem

§ 1. Introduction

Let V be a set of $n (\geq 2)$ points in the Euclidean d -space. The *diameter* of V , $\text{diam}V$, is the maximum of the (Euclidean) distances between points of V . Denote by $e(V)$ the number of pairs $\{x, y\} \subset V$ such that $\|x - y\| = \text{diam}V$, and define $e(d, n)$ to be the maximum of $e(V)$ over all sets V of n points in $\mathbb{R}^d$. We call $V (V \subset \mathbb{R}^d, \#V = n)$ an *extremal configuration* if $n > d$ and $e(V) = e(d, n)$. The case $d = 1$ is not interesting ($e(1, n) = 1$ for $n \geq 2$).

For $d = 2$, it is well known (see [18], pp. 213-214) that $e(2, n) = n$ ($n \geq 3$), and the extremal configurations are fully understood (cf. [29] and [19]). Namely, $V \subset \mathbb{R}^2$ is extremal if and only if $\text{vert}P \subseteq V \subseteq \text{bd}P$ for some Reuleaux polygon P . When P is a Reuleaux k -gon ($3 \leq k \leq n$, k odd), the diameters of V form a self-intersecting k -circuit (the main diagonals of P) plus $n - k$ “dangling edges”. Each dangling edge connects a vertex of P with an interior point of the opposite arc. See Figure 1, where $k = 5$ (a Reuleaux pentagon) and $n = 9$.



– Figure 1 –

If $d = 3$, then $e(3, n) = 2n - 2$ for $n \geq 4$. This has been conjectured by Andrew Vázsonyi (né Endree Weiszfeld⁴, see [10]) and proved in the late 1950's independently by B. Grünbaum, A. Heppes and S. Straszewicz (see [18]). So we refer to the equality $e(3, n) = 2n - 2$, $n \geq 4$, as the *GHS- Theorem*.

The extremal configurations for $d = 3$, however, still form a mostly uncharted territory. Very few examples have been described in the literature⁵. The purpose of this paper is to shed some light on the extremal configurations in dimension 3. We call them the *extremal configurations* and the problem of describing them *the Vázsonyi problem*.

For $d \geq 4$ the quantity $e(d, n)$ grows quadratically with n . More precisely, if $d = 2m$ or $d = 2m + 1$ (d fixed) and $n \rightarrow \infty$, then $e(d, n)$ is asymptotically $(1 - \frac{1}{m}) \cdot \frac{n^2}{2}$; see [18].

It will be convenient to use the notion of diameter graph.

Definition 1.1 (Diameter graph): For $V \subset \mathbb{R}^d$ the geometric graph $D(V) = \langle V, E \rangle$ whose set of vertices is V and whose edges are the pairs $\{x, y\} \subset V$ with $\|x - y\| = \text{diam}V$ is the *diameter graph* of V .

Let us start with a description of the simplest example of a set $V \subset \mathbb{R}^3$ with $\#V = n \geq 4$ and $e(V) = 2n - 2$.

Example 1.1: When $n = 4$, let $V = \{p_0, p_1, p_2, p_3\}$ be the set of vertices of a regular tetrahedron of edge length 1, say. The diameter graph of V is K_4 , and the diameter 1 is attained exactly six ($= 2 \cdot 4 - 2$) times. For $\{i, j, k, l\} = \{0, 1, 2, 3\}$ denote by C_{kl} the circle (of radius $\frac{1}{2}\sqrt{3}$) with center $\frac{1}{2}(p_k + p_l)$ that passes through p_i and p_j , and define A_{ij} to be the closed short circular arc of C_{kl} with endpoints p_i and p_j . Define also $A_{ij}^\circ = A_{ij} \setminus \{p_i, p_j\}$.

⁴Also known as the originator of the “Weiszfeld algorithm”, to approximate successively the point of minimum sum of distances to a finite set of points in $\mathbb{R}^3$ (called the generalized Fermat-Torricelli point).

⁵See [24], Examples 3.1, 3.2, 3.3. Example 3.1 is generalized in our Example 1.1 below; Example 3.2 is the “suspended $(2k - 1)$ -gon” described in Example 1.2 below. Their Example 3.3 of 7 points is brilliant, and as we shall see in another occasion it is obtained from their Example 3.1 for $n = 4$ (4 vertices of a tetrahedron) by an operation called “ball truncation” described in [21]; cf. also Example 9.2 and Remark 9.1 below for the flavor of this operation.

Then A_{ij}° is the locus of all points $x \in \mathbb{R}^3$ that satisfy

$$\|x - p_k\| = \|x - p_l\| = 1, \quad \|x - p_i\| < 1, \quad \|x - p_j\| < 1.$$

For $n > 4$, take V to be the union of $\{p_0, p_1, p_2, p_3\}$ and a set of $n-4$ additional points on some A_{ij}° , say on A_{12}° . The diameter 1 is attained six times at the edges of the tetrahedron, and twice more ($\|x - p_3\| = \|x - p_4\| = 1$) for each additional point x . So far this configuration was known to Grünbaum [13], Heppes [16] and Straszewicz [27] (see also [24], Example 3.1), and here is a generalization of it.

In [26], Lemma 4.2, we have shown in detail that if $x \in A_{ij}^\circ$, $y \in A_{i',j'}^\circ$, then $\|x - y\| < 1$ provided $\{i, j\} \cap \{i', j'\} \neq \emptyset$ (i.e., x and y lie on the same arc or on adjacent arcs), whereas $\|x - y\| > 1$ if $\{i, j\} \cap \{i', j'\} = \emptyset$, i.e., x and y lie on opposite arcs. (A more detailed form of this lemma with a geometric proof is given in Lemma 8.1 below.) Thus we obtain an extremal configuration by taking the vertices p_0, p_1, p_2, p_3 of the tetrahedron, and merely choosing $n-4$ additional points on three mutually adjacent arcs A_{ij}° . These three arcs either share a vertex (like $A_{01}^\circ, A_{02}^\circ, A_{03}^\circ$), or miss a vertex (like $A_{12}^\circ, A_{23}^\circ, A_{31}^\circ$).

Another simple class of extremal configurations can be described as follows.

Example 1.2 (the suspended $(2k-1)$ -gon) (cf. [24], Example 3.2): Let $P = P_{2k-1}$ ($k \geq 2$) be a regular convex $(2k-1)$ -gon of diameter 1 located in the (x, y) -plane of $\mathbb{R}^3$ and with center at the origin o . The diameter of P is attained $2k-1$ times at the main diagonals of P . On the z -axis mark a point c at distance 1 from the vertices of P and put $V = \{c\} \cup \text{vert}P$. Then $n =_{\text{def}} \#V = 2k$, $\text{conv}V$ is a pyramid and the diameter of V is attained $4k-2 = 2n-2$ times: at the main diagonals of the base P , and at the edges emanating from the apex c . Note that the diameter graph $D(V)$ of V is an n -wheel in the sense of graph theory. (An n -wheel is a graph consisting of an $(n-1)$ -circuit plus one universal vertex, the center of the wheel.) For later reference we call this example *a suspended $(2k-1)$ -gon*. When $k \geq 3$, this example allows some perturbation of the points of $V \setminus \{c\}$ on the unit sphere centered at c . A detailed account of this possibility will be given in a subsequent paper.

Remark: The suspended $(2k-1)$ -gon of Example 1.2 serves in Example

9.2 below to construct yet another family of extremal configurations by a construction called “ball truncation”; cf. also Remark 9.1, 2) below.

By these examples one sees that $e(3, n) \geq 2n - 2$ for $n \geq 4$.

The existing proofs that $e(3, n) \leq 2n - 2$ for $n \geq 4$ proceed by induction on n . The case $n = 4$ is obvious. If $\#V = n + 1$ and a point $x \in V$ is incident with at most two diameters (i.e., at most two points of V are at distance $\text{diam}V$ from x), then we apply the induction hypothesis to $V \setminus \{x\}$ and find that $e(V) \leq (2n - 2) + 2 = 2(n + 1) - 2$. Thus the only case that needs some extra consideration is when each point $x \in V$ is at distance $\text{diam}V$ from at least three other points of V . This “extra consideration” consists of three steps:

1. Construction of a convex body $\mathcal{B}(V)$ (“ $\mathcal{B}$ ” stands for “Ball”) related to V . In fact, $\mathcal{B}(V)$ is defined as the intersection of all balls of radius $\text{diam}V$ with centers in V .
2. Defining a “spherical face structure” on the boundary of $\mathcal{B}(V)$ and establishing a relation between this face structure and the diameter graph $D(V)$ of V .
3. Applying Euler’s polyhedral formula ($v - e + f = 2$) to the face structure of $\mathcal{B}(V)$, to conclude that $e(V) \leq 2n - 2$.

We follow these steps carefully and conclude that if $e(V) = 2n - 2$, then the face structure of $\mathcal{B}(V)$ is *strongly self-dual*, i.e., it admits a self-duality of order 2 (involution) which is strong in the following sense: it is fixed-point free when acting as an automorphism of the first barycentric subdivision of the boundary cell complex of $\mathcal{B}(V)$. Nevertheless, the face structure of $\mathcal{B}(V)$, i.e., the poset of its faces, may differ considerably from that of an ordinary convex 3-polytope. The intersection of two facets may have more than one connectivity component, and there may be digonal facets. Consequently the set of faces, partially ordered by inclusion, need not be a lattice⁶, and its

⁶So it is *mistakenly* called a “face lattice” systematically in [3], p. 256, lines 21, 29, 36, p. 260, line 15, and in [2], p. 211, lines 24, 30. In [2], Corollary 6.8 (p. 213), it is shown that, under a strong assumption on V explained in footnote 15 below, $\mathcal{B}(V)$ is a lattice indeed, but this assumption takes the discussion far from the generic case.

1-skeleton forms a planar graph that is not necessarily 3-connected. This is true even if V is a critical configuration (cf. Proposition and Definition 2.1 below). There is a critical configuration $V \subset \mathbb{R}^3$, $\#V = 8$, for which $\mathcal{B}(V)$ is not combinatorially equivalent to a 3-polytope. The configuration will be described in a subsequent paper. There we shall construct new examples of extremal configurations that are modelled after some types of strongly self-dual convex 3-polytopes.

Notation: Our considerations take place in Euclidean d -dimensional space $\mathbb{R}^d$, $d \geq 2$, with origin o , mainly referring to the case $d = 3$. We abbreviate, as usual, aff, conv, vert, int, relint, bd, relbd for *affine hull*, *convex hull*, *vertex set*, *interior*, *relative interior*, *boundary*, and *relative boundary*, respectively.

For $p \in \mathbb{R}^3$ define by $B(p) =_{\text{def}} \{x \in \mathbb{R}^3 : \|x - p\| \leq 1\} = p + B(o)$ the *solid ball centered at p* , where $\|\cdot\|$ is the usual Euclidean norm in $\mathbb{R}^3$, and by $S(p) =_{\text{def}} \text{bd}B(p) = \{x \in \mathbb{R}^3 : \|x - p\| = 1\}$ the *unit sphere centered at p* , usually abbreviated S_p .

The *circumball* of a bounded set $W \subset \mathbb{R}^3$ is the (unique) closed ball of smallest radius that includes W . We denote by $\text{cr}(W)$ (read: circumradius of W), resp. $\text{cc}(W)$ (read: circumcenter of W), the radius, resp. center, of the circumball, and define $\text{cr}(W) = 0$ when $\#W = 1$, and $\text{cr}(W) = \infty$ when W is unbounded. (When $\text{cr}(W) = \infty$, $\text{cc}(W)$ can be taken as any point of $\mathbb{R}^3$.)

§ 2. Critical configurations

Let V be an extremal configuration for the Vázsonyi problem with n points, $n \geq 4$. If $n = 4$, then the diameter graph $D(V)$ is just K_4 . If $n \geq 4$, choose a point $v \in V$ of minimal valence, say δ , in $D(V)$. Then $e(V \setminus \{v\}) = e(V) - \delta = 2n - 2 - \delta$. Note that $\text{diam}(V \setminus \{v\}) = \text{diam}V$, since $\delta \leq n - 1 < 2n - 2$, and thus the removal of v does not obliterate all diameters of V . Assuming the GHS-Theorem (an extended version of which is proved in Theorem 7.1 below), $e(V \setminus \{v\}) \leq 2(n - 1) - 2$. It follows that $\delta \geq 2$. Moreover, if $\delta = 2$, then $V \setminus \{v\}$ is again extremal. This leads us to the following

Proposition – Definition 2.1 (Critical Configuration): *Every point of*

*an extremal configuration (for the Vázsonyi problem) is incident with at least two diameters. If every point of an extremal configuration is incident with at least three diameters, then V is **critical**.*

Examples: The suspended $(2k-1)$ -gon (Example 2.1 above) yields a critical configuration. The 7-point set described in [24], Example 3.3 (mentioned above), is critical as well.

Corollary 2.1: *The diameter graph $D(V)$ of an extremal configuration V does not contain two adjacent 2-valent vertices.*

Proof: If v and w are adjacent 2-valent vertices of $D(V)$, the removal of v leaves w as a 1-valent vertex of $D(V \setminus \{v\})$, and thus $V \setminus \{v\}$ is not extremal, by Proposition 2.1. So $D(V \setminus \{v\})$ has less than $2\#(V \setminus \{v\}) - 2$ diameters. Hence $D(V)$ has less than $2\#(V \setminus \{v\}) - 2 + 2 = 2\#V - 2$ diameters, i.e., V is not extremal, a contradiction. ■

If an extremal configuration V (with $\#V \geq 5$) is not critical, then successive removal of 2-valent vertices $v_1, v_2, \dots, v_k$ (the vertex v_i being 2-valent in $D(V \setminus \{v_1, v_2, \dots, v_{i-1}\})$) will finally produce a critical subconfiguration V_k with $n - k$ points. This will happen after at most $n - k$ steps. Note that the sequence $v_1, v_2, \dots, v_k$ of removed vertices is not necessarily unique. We shall see, however, that the final critical subconfiguration is unique. Denote the set V_k by C (C stands for “critical”). This set C has the following properties:

- 1) $C \subseteq V$, and C is extremal,
- 2) $D(C)$ is a spanned subgraph of $D(V)$.

This follows from the fact, mentioned above, that removal of one point from an extremal configuration does not affect its diameter, and that removal of a 2-valent vertex from an extremal configuration leaves it extremal.

- 3) Every vertex of $D(C)$ has valence ≥ 3 .
- 4) For any set V' , $C \subsetneq V' \subseteq V$, the graph $D(V')$ has a vertex of valence ≤ 2 .

In fact, if v_i is the first vertex in the sequence $v_1, v_2, \dots, v_k$ of removed vertices that belongs to V' , then the valence of v_i in $D(V')$ is smaller than or equal to the valence of v_i in $D(V \setminus \{v_1, v_2, \dots, v_{i-1}\})$, which is 2.

Next we show that C is uniquely determined by the properties 1) – 4) above in the following strong sense.

Theorem 2.1: *If C' is a critical subconfiguration of V , then $C' \subseteq C$.*

Proof: Assume, by r.a.a., that C' is a critical subconfiguration of V and $C' \not\subseteq C$. This means that $C \cup C' \not\subseteq C$. Now consider property 4), with $V' = C \cup C'$. There is a vertex $x \in C \cup C'$ whose valence in $D(C \cup C')$ is ≤ 2 . This is impossible, since if $x \in C$ or $x \in C'$, then the valence of x in $D(C)$ or in $D(C')$, respectively, is already ≥ 3 . ■

Corollary 2.2: *The set C is the unique maximal critical subconfiguration of V , where “maximal” here is with respect to inclusion as well as to cardinality.*

The question whether a critical configuration may still include an extremal configuration as a proper subset (and therefore includes a smaller critical configuration) turns out to be delicate. The answer is (surprisingly) “yes”. The above mentioned example $V \subset \mathbb{R}^3, \#V = 8, \text{diam} V = 1$, for which $\mathcal{B}(V)$ is not 3-polytopal, is critical and has an extremal subconfiguration of four points. Meanwhile we just introduce the relevant technical term.

Definition 2.2: An extremal configuration V is *strongly critical* if no proper subset of V is extremal.

We conjecture that an extremal configuration is strongly critical iff the spherical face structure of its Ball polytope is isomorphic to that of an ordinary 3-polytope; cf. Conjecture 9.2 below.

The only critical configurations we have seen so far are the suspensions of regular $(2k-1)$ -gons ($k \geq 2$), whose diameter graph is a $2k$ -wheel. For $k = 2$, these include the set of vertices of a regular tetrahedron. In the planned subsequent paper we shall construct a large variety of critical configurations, with an arbitrarily large number of points. Most of the rest of this paper is devoted to the study of Ball polytopes and their relation to the extremal configurations. A detailed account of this confronts us with the notions of spindle convexity and spherical convexity discussed in paragraphs 3 and 4 below.

§ 3. Spindle convexity, ball hulls and the Ball connection

In this section we introduce the notion of spindle convexity and the operations $S \rightarrow \mathfrak{B}(S)$ (Ball hull) and $S \rightarrow \mathcal{B}(S)$ (Ball set) for subsets S of $\mathbb{R}^3$, and establish the basic properties of these notions and their interrelations.

Definition and Notation 3.1.: A set $S \subset \mathbb{R}^3$ is *spindle-convex* (abbr. *s-convex*) if for any two distinct points $a, b \in S$, and for any circle C of radius $r \geq 1$ that passes through a and b , the short circular arc of C with endpoints a, b is also included in S . We denote by $sp(a, b)$ (read: spindle a, b) the union of these arcs including the segment $[a, b]$.

Remarks 3.1:

1. If $S \subset \mathbb{R}^3$ is *s-convex*, then S is strictly convex. The proof is easy and left to the reader, with just one hint: If S is *s-convex*, $a, b \in S$ and $y \in \text{relint}[a, b]$, then y is contained in a short circular arc of radius $r \geq 1$ whose two endpoints, x, z say, belong to $sp(a, b) \setminus [a, b] \subset S$, and by the *s-convexity* of S we have $y \in sp(x, z) \setminus [x, z] \subset S$.
2. If $\|a - b\| < 2$, then the union of all short circular arcs of radius $r \geq 1$ with endpoints a and b , including the segment $[a, b]$, forms a spindle with endpoints a and b . When $\|a - b\| \geq 2$, the union of these short circular arcs, including semicircles with endpoints a and b , forms a solid ball centered at $c = \frac{1}{2}(a + b)$ with radius $\frac{1}{2}\|a - b\|^7$. In both cases we denote the spindle with endpoints a and b by $sp(a, b)$.
3. Spindle convexity was studied under the name “Überkonvexität” by Mayer [23]. We introduced this notion in [21]; cf. also [2], p. 203.

An immediate consequence of this definition is

Proposition 3.1: *The intersection of any family of s-convex sets in $\mathbb{R}^3$ is s-convex.*

⁷So it is *not* $\mathbb{R}^3$ as stated in [2], p. 203, line 11.

Lemma 3.1 (*s*-convexity of the unit ball): *A closed ball of radius 1 in $\mathbb{R}^3$ is *s*-convex.*

Lemma 3.1 will follow from a refined version, namely Lemma 3.1' below. We will need this refined version in the sequel (e.g., in the proof of Proposition 5.1 (i)).

Lemma 3.1' (simplified Sallee Lemma): *Let $B = B(z, 1) \subset \mathbb{R}^3$ be a closed unit ball centered at z . Let a, b be two distinct points in B . Let C be a circle of radius $r \geq 1$ with center c that passes through a and b . Let $A \subset C$ be the open short circular arc with endpoints a, b . (When a, b are opposite points of C , A denotes any of the two open semicircles of C with endpoints a, b .) Then $A \subset \text{int}B$, with just one exception: If $a, b \in \partial B$ and $r = 1$, then $c = z$ and $C \subset \partial B$ ⁸.*

Proof: Put $H =_{\text{def}} \text{aff}C$ and $C' =_{\text{def}} H \cap \partial B$, $D' =_{\text{def}} H \cap B = \text{conv}C'$. C' is a circle of radius $r' \leq 1$, with center c' . Note that $r' < 1$, unless H passes through the center z of B , in which case $r' = 1$ and $c' = z$. Now consider the circle C (with radius $r \geq 1$ and center c) and the circle C' (with radius $r' \leq 1$ and center c'). Both these circles lie in H .

If $a \in \text{int}B$ or $b \in \text{int}B$, then C meets $\text{relint}D'$. But $C \not\subseteq D'$, since $r \geq r'$, hence C meets C' at two points, which divide C into two unequal open arcs. The short arc is contained in $\text{relint}D'$ and includes $A(= \widehat{ab})$, and the long one is exterior to $\text{relint}D'$. The same holds true if both a and b are boundary points of B (i.e., $a, b \in C \cap C'$) provided $r > r'$, and even when $r = r'(= 1)$ provided $C \neq C'$ (i.e., $c \neq c'$). There remains the case where $a, b \in \partial B$, $r' = r = 1$ (hence $c' = z$), and $c = c'$. In this case $C' = C$, and thus $A \subset \partial B$. ■

Corollary 3.1.: *The intersection of any family of closed unit balls in $\mathbb{R}^3$ is *s*-convex.*

Under closedness assumption the converse statement of Corollary 3.1 holds as well; see Theorem 3.1 below.

⁸This lemma goes back essentially to Sallee in [25], Lemma 3.3, (p. 317). The lemma appears there in a rather distorted form, which is repeated in [3], Lemma 1.1 (SL1). This last formulation also lacks the delicate exceptional case $a, b \in \partial B$ and $r = 1$ appearing in our formulation. Some overlaps between [25], [2], [3], and the present paper do exist, and we will point out on them occasionally in footnotes.

Definition 3.2 (Ball hull): For a set $S \subset \mathbb{R}^3$, the intersection of all closed unit balls in $\mathbb{R}^3$ that include S , denoted by $\mathfrak{B}(S)$, is the *Ball hull* of S .

Remarks: It follows that $\mathfrak{B}(\emptyset) = \emptyset$ and $S \subset T \implies \mathfrak{B}(S) \subset \mathfrak{B}(T)$. If the circumradius $\text{cr}(S)$ of S is > 1 , then S is not included in any unit ball, and $\mathfrak{B}(S) = \mathbb{R}^3$ (= the intersection of the empty family of unit balls in $\mathbb{R}^3$). If $\text{cr}(S) = 1$, then $\mathfrak{B}(S)$ is just the unique closed unit ball that includes S .

In all cases $S \subset \mathfrak{B}(S)$ and $\mathfrak{B}(S)$ is s -convex. Note that $S = \mathfrak{B}(S)$ iff S is the intersection of some family of closed unit balls in $\mathbb{R}^3$. Next we show that every s -convex set in $\mathbb{R}^3$ is the intersection of a family of closed unit balls, i.e., S is s -convex iff $S = \mathfrak{B}(S)$.

Theorem 3.1: *Suppose that S is a closed subset of $\mathbb{R}^3$. Then S is s -convex iff S is the intersection of a family of (closed) unit balls in $\mathbb{R}^3$.*

Remarks 3.2:

1. The family may be empty, in which case $S = \mathbb{R}^3$. So an alternative formulation of Theorem 3.1 is: Suppose $S \subset \mathbb{R}^3$ is closed. S is s -convex iff either $S = \mathbb{R}^3$ or S is the intersection of a nonempty family of unit balls in $\mathbb{R}^3$.
2. This theorem is the s -convexity analogue of the theorem from linear convexity saying that a closed set $S \subset \mathbb{R}^3$ is convex iff either $S = \mathbb{R}^3$ or S is the intersection of a nonempty family of closed half-spaces.

Proof of Theorem 3.1: The “if” part follows from Corollary 3.1 above. To establish the “only if” part, it suffices to prove

Proposition 3.3 (“separation out” by a unit sphere): *Suppose S is a closed, s -convex proper subset of $\mathbb{R}^3$. If $z \in \mathbb{R}^3 \setminus S$, then there is a closed unit ball $B = B_z \subset \mathbb{R}^3$ that includes S and misses z .*

Indeed, if Proposition 3.3 holds true, then $S = \bigcap \{B_z : z \in \mathbb{R}^3 \setminus S\}$. (Note that the family $\{B_z : z \in \mathbb{R}^3 \setminus S\}$ is empty iff $S = \mathbb{R}^3$, which is compatible

⁹The easier “if” part of this theorem seems to be overlooked until now, and the heavier “only if” part seems to appear already in [23]. We formulated the “only if” part in [21], Theorem 1, and it also appears in [2], Lemma 3.1, with a small gap in the proof. All these appearances are independent of each other.


with the equality $S = \cap \{B_z : z \in \emptyset\}$, since the intersection of an empty set of unit balls is $\mathbb{R}^3$.)

Proof of Proposition 3.3: For $S = \emptyset$ the statement is clear. Assume, therefore, that $S \neq \emptyset$ and denote by a the (unique) point of S nearest to z . Let $u = \frac{a-z}{\|a-z\|}$, and let $H \subset \mathbb{R}^3$ be the plane through a perpendicular to u , i.e., $H = \{x \in \mathbb{R}^3 : \langle u, x \rangle = \langle u, a \rangle\}$. Let H^- be the closed half-space bounded by H that contains z , and let H^+ be the other closed half-space bounded by H . It follows from the convexity of S that $S \subset H^+$.

Note that $a \in S$ and S is s -convex, hence strictly convex, and therefore $S \cap H = \{a\}$, i.e., $S \subset \{a\} \cup \text{int} H^+$. The set $\{a\} \cup \text{int} H^+$ is the union of all spheres tangent to H whose centers lie on the open ray $\{a + \lambda u : \lambda > 0\}$. Define $B_z =_{\text{def}} B(a + u)$, the unit ball centered at $a + u$. Clearly $z \notin B_z$. It remains to show that $S \subset B_z$.

Assume, to the contrary, that there is a point $b \in S \setminus B_z$. Clearly $b \neq a$. There is a unique $\rho > 1$ such that b lies on the sphere $S_\rho =_{\text{def}} S^2(a + \rho u, \rho)$ of radius ρ centered at $a + \rho u$. (An explicit calculation shows that $\rho = \frac{\|a-b\|^2}{2\langle b-a, u \rangle}$.)

Let $A \subset S_\rho$ be a short (closed) circular arc of radius ρ with endpoints a, b .



– Figure 2 –

(A is uniquely determined, unless $b = a + 2\rho u$, in which case A is half a great circle (a meridian) on S_ρ .) Let b' be a point on A such that $0 < \|b' - a\| < 2$,

and let A' be the part of A between a and b' . In the plane $\text{aff}A'$ draw a short circular arc A'' of radius 1 with endpoints a and b' , on the same side of the line $\text{aff}(a, b')$ as A' . Since $A' (\subset H^+)$ is tangent to H at a , and A'' is more curved ($1 < \rho$), A'' protrudes into H^- near a ; see Figure 2. We have $A' \subset A \subset S$, hence $b' \in S$ and $A'' \subset S$, since S is s -convex. But $S \subset H^+$, a contradiction. ■

Remark: Of course, the characterization of s -convex sets given in Theorem 3.1 holds true in $\mathbb{R}^d$ for $d \geq 2$.

Definition 3.3 (Ball set): For a subset V of $\mathbb{R}^3$, the set $\mathcal{B}(V) =_{\text{def}} \{y \in \mathbb{R}^3 : (\forall x \in V)[\|x - y\| \leq 1]\}$ is the *Ball set* (or *Ball connection*) of V . Here $\|\cdot\|$ denotes the usual Euclidean norm in 3-space¹⁰.

We can view $\mathcal{B}(V)$ as the intersection of all (closed) unit balls centered at points of V or, alternatively, as the set of centers of all unit balls that include V . Thus we have the following properties.

- (A) $\mathcal{B}(V) = \bigcap \{v + B : v \in V\}$, and
- (B) $\mathcal{B}(V) = \{z \in \mathbb{R}^3 : V \subset z + B\}$, where $B =_{\text{def}} \{x \in \mathbb{R}^3 : \|x\| \leq 1\}$.

Here are some more properties of the operation $V \rightarrow \mathcal{B}(V)$:

- (C) $\mathcal{B}(\emptyset) = \mathbb{R}^3$ (= the intersection of the empty set of unit balls is $\mathbb{R}^3$).
- (D) $V \neq \emptyset \Rightarrow \mathcal{B}(V)$ is compact and s -convex, hence strictly convex (see (A)).
- (E) For $p \in \mathbb{R}^3$, $\mathcal{B}(\{p\}) = p + B = B(p)$ is the unit ball centered at p .
- (F) $\text{cr}(V) = 1 \Rightarrow \mathcal{B}(V) = \text{cc}(V)$, where $\text{cc}(V)$ is the circumcenter of V (i.e., $\mathcal{B}(V)$ is 0-dimensional).
- (G) (i) When $\text{cr}(V) = 1 - \varepsilon < 1$, $\mathcal{B}(V)$ includes a ball of radius $\varepsilon (> 0)$ centered at $\text{cc}(V)$, i.e., $\mathcal{B}(V)$ is full-dimensional.

¹⁰Ball sets in general and Ball polytopes in particular (cf. Definition 3.4 below) were studied by Sallee in [25] in connection with bodies of constant width (cf. footnote 20 (b) below) and, in a more general setting, in [21], [3], and [2].

- (ii) $\text{cr}(V) > 1 \Rightarrow \mathcal{B}(V) = \emptyset$. Hence
- (iii) $\mathcal{B}(V)$ is full dimensional $\Leftrightarrow \text{cr}(V) < 1$ ¹¹.
- (H) $V_1 \subset V_2 \Rightarrow \mathcal{B}(V_2) \subset \mathcal{B}(V_1)$ (inclusion reversion, but strict inclusion does not imply strict reversed inclusion).
- (I) $\mathcal{B}(V) = \mathcal{B}(\text{cl conv} V)$. In fact, $\mathcal{B}(V) = \mathcal{B}(\mathfrak{B}(V))$, since V and $\mathfrak{B}(V)$ are included in exactly the same unit balls.
- (J) $\mathfrak{B}(V)$ is the largest set $X \subset \mathbb{R}^3$ such that $\mathcal{B}(X) = \mathcal{B}(V)$. In fact, if $Y \subset \mathbb{R}^3$ and $Y \not\subset \mathfrak{B}(V)$, then there is a point $y \in Y \setminus \mathfrak{B}(V)$, and there is a point z such that $V \subset z + B$ but $y \notin z + B$. Thus $z \in \mathcal{B}(V)$, but $z \notin \mathcal{B}(Y)$, i.e., $\mathcal{B}(Y) \not\supset \mathcal{B}(V)$.
- (K) $\text{diam} V \leq 1 \Leftrightarrow V \subset \mathcal{B}(V)$.
- (L) $\text{diam} V \leq 1 \Rightarrow \mathcal{B}(V) = \{x \in \mathbb{R}^3 : \text{diam}(V \cup \{x\}) \leq 1\}$.
- (M) (i) $\mathcal{B}(\mathcal{B}(V)) = \mathfrak{B}(V)$ and, in particular,
- (ii) $\mathcal{B}(\mathcal{B}(V)) \supset V$.

Proof: (i) $x \in \mathcal{B}(\mathcal{B}(V))$
 $\Leftrightarrow (\forall y \in \mathbb{R}^3) [(\forall v \in V) (\|y - v\| \leq 1) \Rightarrow (\|x - y\| \leq 1)]$
 $\Leftrightarrow (\forall y \in \mathbb{R}^3) [(V \subset B(y)) \Rightarrow (x \in B(y))] \Leftrightarrow x \in \mathfrak{B}(V)$.

(ii) Follows from $V \subset \mathfrak{B}(V)$ and (i). ■

(N) $\mathcal{B}(\mathcal{B}(\mathcal{B}(V))) = \mathcal{B}(V)$.

To prove (N) note that the inclusion “ $\supset$ ” follows from (M) (ii) above, where V is replaced by $\mathcal{B}(V)$, and “ $\subset$ ” follows again from (M) (ii) by using (H) (inclusion reversion). (Alternatively, use (M) (i) and $\mathcal{B}(V) = \mathcal{B}(\mathfrak{B}(V))$ proved in (I) above.) From (N) we infer:

Corollary: *The operation $V \rightarrow \mathcal{B}(V)$ is injective (i.e., one to one) on its image set, that is, $\mathcal{B}(S) \neq \mathcal{B}(T) \Rightarrow \mathcal{B}(\mathcal{B}(S)) \neq \mathcal{B}(\mathcal{B}(T))$.*

¹¹This simple condition, equivalent to full dimensionality of $\mathcal{B}(V)$, seems to be new. Sallee, e.g., assumes in [25], p. 316, that $\text{diam} V \leq 1$ (in order to ensure full dimensionality of $\mathcal{B}(V)$), which implies $\text{cr}(V) < 1$; see Theorem 5.2 a) below. Thus Sallee’s condition is sufficient, and clearly not necessary for full dimensionality of $\mathcal{B}(V)$.

(O) $\mathcal{B}(V) = V \Leftrightarrow V$ is a set of constant width 1. (Hence, a set $V \subset \mathbb{R}^3$ of constant width 1 is s -convex.)

This characterization of bodies of constant width (through their Ball connection) is known as “the spherical intersection property”, proved, e.g., by Eggleston in [8], Theorem 52 in p. 123, and in [9], pp. 166-167. See also [6], pp. 62-63, for a proof of this characterization. (A related notion is studied in [20].)

Definition 3.4 (Ball polytope): For $V \subset \mathbb{R}^3$ finite and non-empty with $\text{cr}(V) < 1$, the Ball set $\mathcal{B}(V)$ is the *Ball polytope* associated with V .

Note that $\text{cr}(V) < 1 \Leftrightarrow \mathcal{B}(V)$ is full-dimensional (see (G) (iii) above).

Since a Ball polytope is a finite intersection of balls, it is strictly convex (provided $\text{cr}(V) < 1$), i.e., $(\forall x, y \in \mathcal{B}(V)) [x \neq y \Rightarrow \frac{1}{2}(x + y) \in \text{int } \mathcal{B}(V)]$. The Ball polytope $\mathcal{B}(V)$ of a finite set $V \subset \mathbb{R}^3$ (with, say, $\#V > 1$ and $\text{cr}(V) < 1$) has a natural “spherical” face structure denoted by $\mathcal{SF}(\mathcal{B}(V))$. We shall investigate this face structure thoroughly, and show (in § 8–9 below) that for extremal configurations V of the Vázsonyi problem it admits a canonical fixed-point free involutory self-duality. This means an anti-isomorphism (antimorphism, for short) of $\mathcal{SF}(\mathcal{B}(V))$ of order 2 which, when applied to the first barycentric subdivision of $\mathcal{SF}(\mathcal{B}(V))$, acts as a fixed-point free automorphism. (Note that the first barycentric subdivision of $\mathcal{SF}(\mathcal{B}(V))$ is a purely 2-dimensional simplicial complex which is order-isomorphic to the collection of all chains (“flags” in the current terminology) in $\mathcal{SF}(\mathcal{B}(V))$, partially ordered by inclusion.)

The exposition of the spherical face structure $\mathcal{SF}(\mathcal{B}(V))$ of $\mathcal{B}(V)$ can be viewed as a detailed account of the proof of Vázsonyi’s conjecture mentioned in the introduction. The insight gained by this exposition will lead to the construction of many new types of critical configurations in the planned subsequent paper. In their original proof of the Vázsonyi conjecture, Grünbaum, Heppes and Straszewicz (independently) defined the Ball polytope $\mathcal{B}(V)$ for critical V only (they did not use the term “Ball polytope”) and treated its face structure very sketchy. Our more general definition of $\mathcal{B}(V)$ (for general (finite) configurations $V \subset \mathbb{R}^3$) is aimed to gain a better understanding of this face structure. The next paragraph collects basic results on spherical convexity and apexed cones, also needed in the sequel.

§ 4. Spherical convexity and apexed cones

Let $S(o)$ be the unit sphere centered at the origin.

Definition 4.1 (spherical convexity, strict spherical convexity, spherical convex hull, small cap): A set $F \subset S(o)$ is *spherically convex* [*strictly spherically convex*] if for every two distinct non-antipodal points $a, b \in F$ the geodetic line that connects a and b on $S(o)$ (i.e., the short circular arc with endpoints a, b of the unit circle centered at o which passes through a and b) is contained in F [is contained in a relint F , with possible exception of the endpoints a and b]¹².

Examples 4.1: Here is a list of some spherically convex subsets of $S(o)$.

- a) The empty set and any one-point set.
- b) A set of two antipodal points.
- c) $S(o)$ is strictly spherically convex.
- d) Let $a, b \in S(o)$, $a \neq -b$ (i.e., a, b are not antipodal) and let $\text{arc}(a, b)$ be the geodetic line between a and b on $S(o)$. Then $\text{arc}(a, b)$, $\text{relint } \text{arc}(a, b)$, $\text{arc}(a, b) \setminus \{a\}$, and $\text{arc}(a, b) \setminus \{b\}$ are spherically convex. If $a, b \in S(o)$ are antipodal points, then any closed/open/half-open half great circle on $S(o)$ with endpoints a, b is spherically convex.
- e) A great circle of $S(o)$.
- f) A closed hemisphere of $S(o)$ is spherically convex, but not strictly spherically convex.
- g) A small cap of $S(o)$ is strictly spherically convex (see below).
- e) A spherical triangle on $S(o)$ is spherically convex but not strictly spherically convex, since its edges are geodetic arcs.

¹²In [2], Definition 5.3, spherical convexity is defined for sets contained in a hemisphere only. This assumption is not necessary, as we see here, but it is harmless because most spherically convex sets are contained in a hemisphere.

Note that all these examples are the intersection of $S(o)$ with some cone apexed at o . In Corollary 4.1 below it is shown that this situation is typical.

Assume that $V \subset S(o)$. The intersection of all spherically convex sets that contain V (in $S(o)$) is the *spherical convex hull* of V , henceforth denoted by $\text{sph-conv}(V)$. Clearly, $\text{sph-conv}(V)$ is spherically convex and $V \subset W \Rightarrow \text{sph-conv}(V) \subset \text{sph-conv}(W)$.

Let H be a plane that misses the origin o such that $\text{dist}(o, H) \leq 1$. The intersection of $S(o)$ with the closed half-space bounded by H that misses o is a *small cap* of $S(o)$. A small cap is obviously strictly spherically convex.

Definition 4.2 (apexed cone): Let $V \subset \mathbb{R}^3 \setminus \{o\}$ be a compact set. The cone $\text{cone}_o V$ is *apexed* (at o) if there exists a plane $H \subset \mathbb{R}^3$ strictly separating V and o . Equivalently, $\text{cone}_o V$ is apexed (at o) iff there exists a plane H' strictly supporting $\text{cone}_o V$ at o .

Proposition 4.1: *Assume $V \subset S(o)$. Then*

- (i) *if $\text{cone}_o V$ is apexed (at o), then $\text{cone}_o V \cap S(o)$ is contained in a small cap of $S(o)$,*
- (ii) *$\text{cone}_o V \cap S(o)$ is spherically convex in $S(o)$,*
- (iii) *$\text{sph-conv}(V) = \text{cone}_o V \cap S(o)$.*

Proof:

- (i) This follows readily from Definition 4.2.
- (ii) The cases $V = \emptyset$ and $\#V = 1$ are trivial. In the case $\#V = 2$ and if the two points of V are antipodal, $\text{cone}_o V \cap S(o) = V$ and the set of two antipodal points is spherically convex (cf. Examples 4.1 b) above). Let $a, b \in \text{cone}_o V \cap S(o)$ be two non-antipodal points, and let z be a point on the short arc between a and b on the unit circle centered at o passing through a and b . Then $z \in \text{cone}_o(\{a, b\})$, i.e., $z = \lambda a + \mu b$ for some $\lambda, \mu \geq 0$. The point a , resp. b , is a non-negative combination of a finite subset $\{a_1, \dots, a_m\} \subset V$, resp. $\{b_1, \dots, b_l\} \subset V$, i.e., $a = \sum_{i=1}^m \lambda_i a_i$,

resp. $b = \sum_{i=1}^l \mu_i b_i$, for some $\lambda_i \geq 0$ ($1 \leq i \leq m$), resp. $\mu_i \geq 0$ ($1 \leq i \leq l$).
Thus

$$z = \lambda a + \mu b = \lambda \sum_{i=1}^m \lambda_i a_i + \mu \sum_{i=1}^l \mu_i b_i = \sum_{i=1}^m \lambda \lambda_i a_i + \sum_{i=1}^l \mu \mu_i b_i,$$

which is a non-negative combination of $\{a_i : 1 \leq i \leq m\} \cup \{b_i : 1 \leq i \leq l\} \subset V$, hence $z \in \text{cone}_o V$. Since $\|z\| = 1$, $z \in \text{cone}_o V \cap S(o)$.

- (iii) By (ii), $\text{sph-conv} V \subset \text{cone}_o V \cap S(o)$. To prove “ $\supset$ ” assume that F is a spherically convex subset of $S(o)$ containing V . We show that $\text{cone}_o V \cap S(o) \subset F$. Let $z \in \text{cone}_o V \cap S(o)$. Then z is a non-negative combination of a finite subset $W \subset V$. By Carathéodory’s Theorem (2-dimensional case) there are three points $w_1, w_2, w_3 \in W$ no two of which are antipodal such that

$$z = \lambda_1 w_1 + \lambda_2 w_2 + \lambda_3 w_3, \tag{1}$$

where $\lambda_i \geq 0$ ($1 \leq i \leq 3$), i.e., $z \in \text{cone}_o(\{w_1, w_2, w_3\})$.

If there is a λ_i ($1 \leq i \leq 3$) such that $\lambda_i = 0$, say $\lambda_3 = 0$, then $z \in \text{cone}_o(\{w_1, w_2\})$, i.e., z lies on the short arc connecting w_1 and w_2 of the unit circle with center o passing through w_1 and w_2 . Since $w_1, w_2 \in F$ and F is closed to such short circles $z \in F$.

Assume now that $\lambda_1, \lambda_2, \lambda_3 > 0$. The vector $\lambda_1 w_1 + \lambda_2 w_2$ is a scalar product of the unit vector $w =_{\text{def}} \frac{\lambda_1 w_1 + \lambda_2 w_2}{\|\lambda_1 w_1 + \lambda_2 w_2\|}$ by the positive scalar $\lambda =_{\text{def}} \|\lambda_1 w_1 + \lambda_2 w_2\|$. The unit vector w lies on the short arc with endpoints w_1, w_2 of the unit circle centered at o and passing through w_1 and w_2 , hence $w \in F$. By (1) we have $z = \lambda w + \lambda_3 w_3$, hence z lies on the short arc with endpoints w and w_3 of the unit circle centered at o and passing through w and w_3 , hence $z \in F$. This proves (iii). ■

From Proposition 4.1 (iii) we conclude

Corollary 4.1: *If $V \subset S(o)$ is spherically convex, then $V = \text{cone}_o V \cap S(o)$.*

Theorem 4.1 below is stated just for a general background without proof, since it is not needed in the sequel.

Theorem 4.1: *If $F \subset S(o)$ is a closed, spherically convex set not containing a pair of antipodal points, then*

- a) *F is contained in a small cap of $S(o)$,*
- b) *F is the intersection of a family of closed hemispheres of $S(o)$.*

A close connection between spindle convexity and spherical convexity is enunciated in the following statement (compare [2], Lemma 5.6).

Proposition 4.2: *If $K \subset \mathbb{R}^3$ is spindle convex, then $S(o) \cap K$ is spherically convex on $S(o)$, and either $K = \mathbb{R}^3$ or $K = B(o)$ or $S(o) \cap K$ is contained in small cap of $S(o)$.*

Proof: If $\#(S(o) \cap K) \leq 1$, there is nothing to prove. Assume $\#(S(o) \cap K) \geq 2$, let $a, b \in F =_{\text{def}} S(o) \cap K$, $a \neq b$, and let $\text{arc}(a, b)$ be a short arc (possibly half a circle) of radius 1 on $S(o)$ with endpoints a, b . Since K is s -convex, $\text{arc}(a, b) \subset K$, hence $\text{arc}(a, b) \subset F$ and F is spherically convex. By Remark 3.2, 1) either $K = \mathbb{R}^3$ or K is the intersection of a non-empty family of unit balls. In the second case $K \subset B(p)$ for some $p \in \mathbb{R}^3$ (the unit ball centered at p). If $K = B(o)$, we are done. Otherwise $K \subset B(p)$ for some $p \neq o$, hence $S(o) \cap K \subset S(o) \cap B(p)$, and since $p \neq o$, clearly $S(o) \cap B(p)$ is a small cap of $S(o)$. ■

Remark: The intersection $S(o) \cap K$ alluded to above need not be *strictly* spherically convex. Let $a, b, c \in S(o)$ be three points no two of which are antipodal such that $o \notin \text{aff}\{a, b, c\}$. Then $\Delta =_{\text{def}} \text{sph-conv}(\{a, b, c\})$ is a spherical triangle which is not strictly spherically convex (the edges of Δ are geodesic arcs of $S(o)$). Put $K =_{\text{def}} \cap \{B \subset \mathbb{R}^3 : B \text{ is a unit ball and } \Delta \subset B\}$. K is spindle convex (by Theorem 3.1), and it can be shown that $S(o) \cap K = \Delta$, thus it is not strictly spherically convex.

§ 5. The face structure of Ball polytopes

Essential and inessential points:

Let $V \neq \emptyset$ be a finite set of points in $\mathbb{R}^3$. We assume that $\text{cr}(V) < 1$; ipso facto $\mathcal{B}(V)$ is full-dimensional, and $\text{diam}V < 2$.

Definition 5.1: A point $v \in V$ is *essential* if $\mathcal{B}(V) \subsetneq \mathcal{B}(V \setminus \{v\})$, otherwise (if $\mathcal{B}(V) = \mathcal{B}(V \setminus \{v\})$) it is *inessential*.

Note that v is essential iff there is a unit ball that includes $V \setminus \{v\}$ and misses v . In particular, if v is essential, then v is an extreme point of V . Moreover, this characterization of essential points shows that if v is essential in V , then v is essential in every subset of V that contains v . Denote by $\text{ess}(V)$ the set of essential points of V . Soon we shall show that $\mathcal{B}(\text{ess}(V)) = \mathcal{B}(V)$ (Theorem 5.1 below).

Lemma 5.1: *We have $v \in \text{ess}(V)$ iff there is a point $x \in \mathbb{R}^3$ such that $\|x - v\| = 1$ and $\|x - w\| < 1$ for $w \in V \setminus \{v\}$ ¹³.*

Proof: ($\Rightarrow$) Assume $v \in \text{ess}(V)$. Let c be the center of a unit ball that includes $V \setminus \{v\}$ but misses v , and let b be an interior point of $\mathcal{B}(V)$. Then $\|b - w\| < 1$ for $w \in V$, $\|c - v\| > 1$ and $\|c - w\| \leq 1$ for $w \in V \setminus \{v\}$. For $0 \leq \vartheta \leq 1$, define $a(\vartheta) =_{\text{def}} (1 - \vartheta)b + \vartheta c$. Choose ϑ , $0 < \vartheta < 1$, such that $\|a(\vartheta) - v\| = 1$, and define $x = a(\vartheta)$. Then $\|x - w\| < 1$ for $w \in V \setminus \{v\}$.

($\Leftarrow$) Assume $\|x - v\| = 1$ and $\|x - w\| < 1$ for $w \in V \setminus \{v\}$. Choose a positive number ε such that $\|x - w\| \leq 1 - \varepsilon$ for $w \in V \setminus \{v\}$, and define $x' =_{\text{def}} x + \varepsilon(x - v)$. Then $\|x' - v\| = 1 + \varepsilon > 1$, but $\|x' - w\| \leq 1$ for $w \in V \setminus \{v\}$, hence $v \in \text{ess}(V)$. ■

Next we show that the removal of an inessential point does not change the status (essential/inessential) of the remaining points with respect to the remaining set. We have seen already that if $w \in V \setminus \{v\}$ is essential in V , then it is essential in $V \setminus \{v\}$. The next lemma does the rest.

Lemma 5.2: *Assume $V \subset \mathbb{R}^3$ is finite, with $\text{cr}(V) < 1$. If v and v' are inessential in V , then v' is inessential in $V \setminus \{v\}$.*

Proof: Assume, on the contrary, that $v' \in \text{ess}(V \setminus \{v\})$. By Lemma 5.1 there is a point $x \in \mathbb{R}^3$ such that $\|x - v'\| = 1$, and $\|x - w\| < 1$ for $w \in V \setminus \{v, v'\}$. What about $\|x - v\|$? If $\|x - v\| < 1$, then v' is essential in V , contrary to our hypothesis. If $\|x - v\| > 1$, define $x(\varepsilon) =_{\text{def}} (1 - \varepsilon)x + \varepsilon v'$. For $\varepsilon > 0$ sufficiently

¹³See [3], Proposition 1.1, for a similar statement.

small, we find that $\|x(\varepsilon) - v'\| < 1$, $\|x(\varepsilon) - v\| > 1$, and $\|x(\varepsilon) - w\| < 1$ for all $w \in V \setminus \{v, v'\}$. Thus $v \in \text{ess}(V)$, contrary to our hypothesis. If $\|x - v\| = 1$, then the points x, v, v' form an isosceles triangle with $\|x - v\| = \|x - v'\| = 1$. Move the point x slightly parallel to the base $[v, v']$ of this triangle. The resulting point x' satisfies: $\|x' - v\| > 1$, $\|x' - v'\| < 1$ (or vice versa), and $\|x' - w\| < 1$ for all $w \in V \setminus \{v, v'\}$. Thus $v \in \text{ess}(V)$ (or $v' \in \text{ess}(V)$), contrary to our hypothesis. ■

Theorem 5.1: *Assume that $V \subset \mathbb{R}^3$ is finite with $\text{cr}(V) < 1$. Then $\mathcal{B}(V) = \mathcal{B}(\text{ess}(V))$.*

Proof: Remove the inessential points of V one by one, and apply Lemma 5.2 each time to ensure that all inessential points remain inessential. ■

Clearly, for any proper subset $W \subsetneq \text{ess}(V)$ one has $\mathcal{B}(W) \subsetneq \mathcal{B}(V) (= \mathcal{B}(\text{ess}(V)))$, hence, by Theorem 5.1, $\text{ess}(V)$ is the smallest subset X of V satisfying $\mathcal{B}(X) = \mathcal{B}(V)$. This brings us to

Definition 5.2: A finite set $V \subset \mathbb{R}^3$ satisfying $\text{cr}(V) < 1$ and $V = \text{ess}(V)$ is *tight*¹⁴.

If $V \subset \mathbb{R}^3$ is finite and not tight, then after removal of the inessential points from V there remains a (tight) set whose ball polytope is the same $\mathcal{B}(V)$. Hence, in addition to the standard assumption $\text{cr}(V) < 1$, we will also assume that V is tight when dealing with the face structure of the Ball polytope $\mathcal{B}(V)$ of V . Theorem 5.2 a), b) below guarantee these two properties in case that $V \subset \mathbb{R}^3$ is extremal for the Vázsonyi problem.

Theorem 5.2: *Assume that $V \subset \mathbb{R}^3$ is finite and $\text{diam} V = 1$. Then*

- a) $\text{cr}(V) < 1$.
- b) *If a point $v \in V$ is incident with (at least) two diameters of V , then $v \in \text{ess}(V)$.*
- c) *If V is extremal for the Vázsonyi problem, then V is tight.*

¹⁴In [2], Definition 6.3, such a set is called *reduced*, and in [3], p. 258, it is called *non-redundant*. Sallee in [25] had no name for this notion, though he clearly had it in mind in his (confused) definition of a *finite intersection of balls* in p. 316, lines 1-5.

Proof:

- a) In fact, we have $\text{cr}(V) \leq \sqrt{\frac{3}{8}} (< 1)$, where equality is attained iff V is the set of vertices of a tetrahedron of edge-length 1 (Jung's Theorem; see [8], Theorem 49 in p. 111, or [17]).
- b) Assume $a, b \in V$, $\|a - v\| = \|b - v\| = 1$. Define $c =_{\text{def}} \frac{1}{2}(a + b)$. Denote by H the plane through c perpendicular to $[a, b]$ and by H_a, H_b the closed half-spaces determined by H that contain a and b , respectively. Note that

$$V \subset \mathcal{B}(V) \subset B(a, 1) \cap B(b, 1) = (B(a, 1) \cap H_b) \cup (B(b, 1) \cap H_a). \quad (2)$$

Denote by C the circle with center c that passes through v , i.e., $C = \{x \in \mathbb{R}^3 : \|x - a\| = \|x - b\| = 1\}$. Denote by w the point at distance 1 from v on the ray $\overrightarrow{vc}$. Note that $\|w - x\| < 1$ for every point x on C or inside the disc $\text{conv}C$, except v , i.e., $\text{conv}C \setminus \{v\} \subset \text{int}B(w, 1)$. By spindle convexity of $B(w, 1)$ (Lemma 3.1) we have

$$(B(a, 1) \cap H_b) \cup (B(b, 1) \cap H_a) \subset \{v\} \cup \text{int}B(w, 1). \quad (3)$$

Thus $\|v - w\| = 1$, and by (2) and (3) we have $\|v' - w\| < 1$ for $v' \in V \setminus \{v\}$. This shows, by Lemma 5.1, that v is essential in V .

- c) This follows from a), b) and Proposition-Definition 2.1 above. ■

Now we are ready to introduce the spherical face structure $\mathcal{SF}(\mathcal{B}(V))$ of the Ball polytope $\mathcal{B}(V)$. We assume that V is finite, nonempty with $\text{cr}(V) < 1$, and tight ($V = \text{ess}(V)$). $\mathcal{SF}(\mathcal{B}(V))$ will be a finite 5-tiered poset (not necessarily a lattice) consisting of the empty set $\emptyset$, vertices, edges, facets, and $\mathcal{B}(V)$ itself. We shall introduce and discuss these elements in the following order: First facets, then vertices, then edges.

Facets:

Definition 5.3 (facet of $\mathcal{B}(V)$): For a point $p \in V$ the set $F_p =_{\text{def}} \{x \in \mathcal{B}(V) : \|x - p\| = 1\}$ is a *facet* of $\mathcal{B}(V)$.

By definition, F_p is a compact subset of the unit 2-sphere centered at p , $S_p =_{\text{def}} \{x \in \mathbb{R}^3 : \|x - p\| = 1\}$. Moreover, since V is tight, p is essential in

V , and thus (by Lemma 5.1) there is a point $x \in S_p$ such that $\|x - q\| < 1$ for $q \in V \setminus \{p\}$. The point x is interior to F_p , relative to S_p . Hence $\text{relint} F_p \neq \emptyset$ and $\mathcal{B}(V)$ has exactly $\#V$ facets. For $V = \{p\}$ we have, of course, $F_p = S_p$. When $V \supsetneq \{p\}$, we can write $F_p := S_p \cap \bigcap_{q \in V \setminus \{p\}} \{x \in \mathbb{R}^3 : \|x - q\| \leq \|x - p\|\}$.

For $x \in \mathbb{R}^3$ and $q \in V \setminus \{p\}$ we have $\|x - q\| \leq \|x - p\| \Leftrightarrow \langle x - p, q - p \rangle \geq \frac{1}{2}\|q - p\|^2$. (This can be verified by a straightforward calculation.) The set $H_{p,q} =_{\text{def}} \{x \in \mathbb{R}^3 : \langle x - p, q - p \rangle \geq \frac{1}{2}\|q - p\|^2\}$ is a closed half-space of $\mathbb{R}^3$ that misses p . The intersection of S_p with such a half-space is a small cap of S_p . Thus we have expressed F_p as an intersection of $\#V - 1$ or less small caps of S_p : $F_p = \bigcap_{q \in V \setminus \{p\}} (S_p \cap H_{p,q})$. A small cap $C =_{\text{def}} S_p \cap H_{p,q}$ is strictly spherically convex (in the sense of Definition 4.1 above), hence we have

Corollary 5.2: *If $V \subset \mathbb{R}^3$ is finite, tight, $\#V \geq 2$, and $\text{cr}(V) < 1$, then for $p \in V$ the facet F_p is a strictly spherically convex subset of S_p , contained in a small cap of S_p , compact with a non-empty relative interior $F_p = \{x \in S_p : \|x - q\| \leq 1 \text{ for } q \in V \setminus \{p\}\} = S_p \cap (\mathcal{B}(V \setminus \{p\}))$ and $\text{relint} F_p = \{x \in S_p : \|x - q\| < 1 \text{ for } q \in V \setminus \{p\}\}$.*

Next we show that F_p is topologically a 2-disc (= 2-cell).

Theorem 5.3: *If $F \subset S(o)$ is compact, strictly spherically convex and contained in a small cap of $S(o)$ with $\text{relint} F \neq \emptyset$, then there is a homeomorphism of F to a closed hemisphere of $S(o)$, which maps $\text{relbd} F$ (the relative boundary of F) to a great circle in $S(o)$.*

Proof: Assume, without loss of generality, that the north pole $N(0, 0, 1)$ is relatively interior to F . We notice that for $(\varphi, \vartheta) \in (-\pi, \pi] \times [-\frac{\pi}{2}, \frac{\pi}{2}]$ the vector $(\cos \varphi \cos \vartheta, \sin \varphi \cos \vartheta, \sin \vartheta)$ is a parametric representation of the meridian line of $S(o)$ of fixed longitude $\varphi \in (-\pi, \pi]$ (spherical coordinates).

Define $\vartheta(\varphi) =_{\text{def}} \min \left\{ \vartheta : \begin{pmatrix} \cos \varphi \cos \vartheta \\ \sin \varphi \cos \vartheta \\ \sin \vartheta \end{pmatrix} \in F \right\}$ for $-\pi < \varphi \leq \pi$. Clearly, $-\frac{\pi}{2} < \vartheta(\varphi) < \frac{\pi}{2}$, and one can easily check that $\vartheta(\varphi)$ is a continuous function of φ (this follows from the compactness and the strict spherical convexity of F). A general point of F is of the form $x = (\cos \varphi \cos \vartheta, \sin \varphi \cos \vartheta, \sin \vartheta)$, $\vartheta(\varphi) \leq \vartheta \leq \frac{\pi}{2}$, and $\vartheta = (1 - \lambda)\vartheta(\varphi) + \lambda \cdot \frac{\pi}{2}$, $0 \leq \lambda \leq 1$. Map x to $f(x) =_{\text{def}} (\cos \varphi \cos(\lambda \cdot \frac{\pi}{2}), \sin \varphi \cos(\lambda \cdot \frac{\pi}{2}), \sin(\lambda \cdot \frac{\pi}{2}))$.

This defines a homeomorphism between F and the northern hemisphere of $S(o)$, which maps $\text{relbd}F$ to the equator. $\blacksquare$

A point $x \in \mathcal{B}(V)$ is a boundary point of $\mathcal{B}(V)$ if it satisfies at least one of the inequalities $\|x - p\| \leq 1$ ($p \in V$) as an equality. Thus

$$\text{bd}\mathcal{B}(V) = \bigcup \{F_p : p \in V\}. \quad (4)$$

Theorem 5.4: For $p \in V$, $\text{relbd}F_p = \bigcup \{F_p \cap F_q : q \in V \setminus \{p\}\}$.

Proof: Assume $x \in F_p$. Then $\|x - p\| = 1$ and $\|x - q\| \leq 1$ for $q \in V \setminus \{p\}$. If all these inequalities are strict, then they hold for $x' \in S_p$ instead of x , provided $\|x' - x\| \leq 1 - \|x - q\|$ for $q \in V \setminus \{p\}$.

Thus x is interior to F_p relative to S_p . If, on the other hand, $\|x - q\| = 1$ for some $q \in V \setminus \{p\}$, then the points x, p, q form an isosceles triangle T with $\|x - p\| = \|x - q\| = 1$. (We have $\|p - q\| < 2$, since otherwise $\text{cr}(V) \geq 1$, contrary to our assumptions.) Moving x slightly within $\text{aff}T$ away from q along a unit circle centered at p , we obtain a point $x' \in S_p$ that satisfies $\|x' - q\| > 1$. Hence $x' \notin \mathcal{B}(V)$ and, in particular, $x' \notin F_p$. Thus $x \in \partial F_p$. $\blacksquare$

What can be said about the intersection $F_p \cap F_q$ for $p, q \in V, p \neq q$? By definition $F_p \cap F_q = \{x \in \mathcal{B}(V) : \|x - p\| = \|x - q\| = 1\}$.

The set $C_{pq} =_{\text{def}} \{x \in \mathbb{R}^3 : \|x - p\| = \|x - q\| = 1\}$ is the “equatorial” circle of the lentil shaped body $\mathcal{B}(\{p, q\})$, of radius r_{pq} satisfying $0 < r_{pq} =_{\text{def}} \sqrt{1 - \frac{1}{4}\|p - q\|^2} < 1$ and centered at $c_{pq} =_{\text{def}} \frac{1}{2}(p + q)$, lying on the plane that bisects the segments $[p, q]$ perpendicularly. The intersection of C_{pq} with any unit ball $B(v, 1)$, $v \in V \setminus \{p, q\}$, is a-priori either the whole of C_{pq} , or a closed arc of C_{pq} , or a singleton or empty. It follows that $F_p \cap F_q = C_{pq} \cap \text{bd}\mathcal{B}(V)$ has at most $n - 2$ components (closed arcs or singletons). Next we prove (part (i) of the following proposition) that for $\#V \geq 3$ the intersection $F_p \cap F_q$ is not the whole of C_{pq} , not a singleton, nor empty, which is a key for some later developments.

Proposition 5.1: Assume that $V \subset \mathbb{R}^3$, $\#V \geq 3$ is finite, tight, $\text{cr}(V) < 1$, and let $p, q \in V$. Then

- (i) C_{pq} is a circle of radius $0 < r_{pq} < 1$, and for $v \in V \setminus \{p, q\}$, $C_{pq} \cap B(v, 1)$ is a relatively closed arc of C_{pq} (with two (different) endpoints; ipso facto it has a non-empty relative interior). Equivalently: $\mathcal{B}(v, 1)$ misses a nonempty, proper (relatively open) arc of C_{pq} .
- (ii) $F_p \cap F_q = C_{pq} \cap \mathcal{B}(V)$ is either empty, or it is a finite union of at most $\#V - 2$ closed arcs (connectivity components) of C_{pq} , with each arc having either two endpoints or being a single point.
- (iii) Each endpoint of a connectivity component of $C_{pq} \cap \mathcal{B}(V)$ is common to at least three facets of $\mathcal{B}(V)$ (including F_p and F_q , of course).
- (iv) An isolated point of $C_{pq} \cap \mathcal{B}(V)$ is common to at least four facets of $\mathcal{B}(V)$.

Proof:

- (i) Since V is tight and $\text{cr}(V) < 1$, $\mathcal{B}(\{p, q\}) = \mathcal{B}(p) \cap \mathcal{B}(q)$ is a full-dimensional “lentil” shaped body whose “equator” C_{pq} is a circle of radius r_{pq} , $0 < r_{pq} < 1$. It remains to show that, for $v \in V \setminus \{p, q\}$, $C_{pq} \setminus B(v) \neq \emptyset$ and that $B(v) \cap C_{pq}$ has more than one point. (Note that $B(v, 1)$ is closed, hence $C_{pq} \setminus B(v, 1)$ is relatively open in C_{pq} ; so it cannot be a singleton.)

Assume, by r.a.a., that $C_{pq} \subset B(v)$. The boundary $\text{bd}(B(p) \cap B(q))$ of $B(p) \cap B(q)$ consists of two small caps $\text{cap}(p) =_{\text{def}} S(p) \cap B(q)$ of $S(p)$ and $\text{cap}(q) =_{\text{def}} S(q) \cap B(p)$ of $S(q)$. Let $x \in \text{relint cap}(p)$ ($= \text{cap}(p) \setminus C_{pq}$) and let $C \subset S(p)$ be a great circle of $S(p)$ (of radius 1 and centered at p) passing through x . The intersection $C \cap \text{cap}(p)$ is a short arc of C , both of its endpoints lie on $C_{pq} \subset B(v)$. By Lemma 3.1’ (simplified Sallee Lemma) $\text{relint } A \subset \text{int} B(v)$, unless $v = p$, which is impossible ($v \neq p$ by assumption). Thus $x \in \text{int} B(v)$ and $\text{cap}(p) \subset B(v)$. A similar argument shows that $\text{cap}(q) \subset B(v)$, hence $\text{bd}(B(p) \cap B(q)) = \text{cap}(p) \cup \text{cap}(q) \subset B(v)$. It follows that $B(p) \cap B(q) \subset B(v)$ (by convexity of $B(p) \cap B(q)$), a contradiction to the assumption that v is essential. Thus

$$C_{pq} \not\subset B(v), \tag{5}$$

and similarly (interchanging the roles of q and v) we get

$$C_{pv} \not\subset B(q). \quad (6)$$

By (5) $C_{pq} \cap B(v)$ is either a proper arc of C_{pq} or a singleton or empty. In order to disprove the latter two possibilities we consider two cases.

Case I: $v \in H =_{\text{def}} \text{aff} C_{pq}$.

If $\|v - \frac{1}{2}(p+q)\| \leq r_{pq} (< 1)$, then the circle $B(v) \cap H$ of radius 1 has its center v in $\text{conv} C_{pq}$, it is not contained in $\text{conv} C_{pq}$ (since $r_{pq} \leq 1$), and it does not contain C_{pq} (by (5)), hence it intersects C_{pq} in two points. Assume now that $\|v - \frac{1}{2}(p+q)\| > r_{pq}$, i.e., v lies out of the circle C_{pq} (in the plane H). Then

$$\text{dist}(v, B(p) \cap B(q)) = \text{dist}(v, C_{pq}). \quad (7)$$

Since $\text{cr}\{v, p, q\} < 1$, it follows from (7) that $\text{dist}(v, C_{pq}) < 1$, and by (5) the circle $B(v) \cap H$ intersects C_{pq} in two points.

Case II: $v \notin H$.

Let H^+, H^- be the two open half-spaces bounded by H and assume, without loss of generality, that $\{v, q\} \subset H^-$ and $p \in H^+$.


If $C_{pq} \cap C_{pv}$ is singleton or empty, then – both circles lying on $S(p)$ – either $C_{pv} \subset H \cup H^+$ or $C_{pv} \subset H \cup H^-$. In the first case $C_{pq} \subset B(v)$, contradicting (5), and in the second case $C_{pv} \subset B(q)$, contradicting (6). Thus $\#(C_{pq} \cap C_{pv}) \geq 2$, i.e., $C_{pq} \cap B(v)$ is a proper, non-singleton arc of C_{pq} .

- (ii) By part (i), $F_p \cap F_q = C_{pq} \cap \mathcal{B}(V) = \bigcap \{C_{pq} \cap B(v, 1) : v \in V \setminus \{p, q\}\}$ is a finite intersection of $\#V - 2$ closed arcs on C_{pq} , which is either empty or of the type described in the proposition.
- (iii) Let x be an endpoint of a connectivity component A of $C_{pq} \cap \mathcal{B}(V)$. The point x is common to F_p and F_q , and if it does not belong to any other facet F_v , $v \in V \setminus \{p, q\}$, then $x \in \text{int} \mathcal{B}(V \setminus \{p, q\})$. It follows that there is a whole neighborhood of x on C_{pq} included in $\mathcal{B}(V)$, contradicting the assumption that x is an endpoint of a connectivity component of $C_{pq} \cap \mathcal{B}(V)$.
- (iv) Follows from (i) and (iii). ■

The following example shows that the intersection of two facets $F_p \cap F_q = C_{pq} \cap \mathcal{B}(V)$ can be, indeed, *any* finite disjoint union of closed (not necessarily short) arcs of C_{pq} , including singletons. In the next paragraph we define the 1-skeleton of a ball polytope and refer to this as a basic example of a ball polytope whose 1-skeleton is not 3-connected.

Example 5.1: Let $p = p(0, 0, h)$, $q = q(0, 0, -h)$, $0 < h < 1$, and let $C =_{\text{def}} S(p) \cap S(q) = C_{pq}$ be a circle in the (x, y) -plane centered in the origin of radius $\sqrt{1 - h^2}$, $0 < \sqrt{1 - h^2} < 1$. The set $\text{bd}\mathcal{B}(\{p, q\})$ is the union of two small caps of $S(p)$ and $S(q)$, respectively, the relative boundary of both of which is C . Assume that C is positively oriented (counterclockwise in all our figures).

Notation: For $u, v \in C$, $u \neq v$, denote by $\text{orarc}(u, v)$ the positively oriented arc of C beginning in u and terminating in v . ($\text{orarc}(u, v)$ may be a short as well as a long arc of C , of course).



– Figures 3 and 4 –

Mark successively n points $p_0, p_1, \dots, p_{n-1}$, $n \geq 2$, on C so that the cyclic order of the indices $0, 1, \dots, n-1$ is compatible with the positive orientation on C , i.e., for $1 \leq i \leq n$ the arc $\text{orarc}(p_{i-1}, p_i)$ (the indices are taken modulo n , so that $p_n = p_0$) does not contain any point p_j ($1 \leq j \leq n$) in its relative interior. For $1 \leq i \leq n$ choose a point v_i in the (x, y) -plane such that

- i) $\|v_i - p_{i-1}\| = \|v_i - p_i\| = 1$, and
- (ii) $\|v_i - x\| > 1$ for $x \in \text{relint}(\text{orarc}(p_{i-1}, p_i))$.

Conditions (i) and (ii) uniquely determine v_i , and they imply that v_i and $\text{orarc}(p_{i-1}, p_i)$ lie on opposite sides of the chord $[p_{i-1}, p_i]$ of C ; see Figures 3 and 4.

Thus the unit ball $B(v_i)$ misses $\text{relint}(\text{orarc}(p_{i-1}, p_i))$ and includes its complementary arc on C , namely $\text{orarc}(p_i, p_{i-1})$.

Choose any subset $W \subseteq \{v_0, v_1, \dots, v_{n-1}\}$ and define $V =_{\text{def}} \{p, q\} \cup W$. Then $\text{cr}(V) < 1$, V is tight, and

$$\begin{aligned} F_p \cap F_q = C_{pq} \cap \mathcal{B}(V) = (\bigcup \{ \text{orarc}(p_{i-1}, p_i) : \\ v_i \notin W, 1 \leq i \leq n \} \cup \{ p_i : v_i, v_{i+1} \in W, 1 \leq i \leq n \} . \end{aligned} \quad (8)$$

Hence, by choosing W at will, we have full control of which arc $\text{orarc}(p_{i-1}, p_i)$ of C belongs to $F_p \cap F_q = C \cap \mathcal{B}(V)$, and of which not. E.g., if $W = \{v_0, v_1, \dots, v_{n-1}\}$, then $F_p \cap F_q = \{p_0, p_1, \dots, p_{n-1}\}$ because $B(v_i)$ misses $\text{relint}(\text{orarc}(p_{i-1}, p_i))$ for $1 \leq i \leq n$.

Note that if $v_i \in W \subset V$, then the facet F_{v_i} of $\mathcal{B}(V)$ has exactly two “edges”, both of which are short arcs with endpoints p_{i-1}, p_i of the circles $C_{pv_i} = S(p) \cap S(v_i)$ and $C_{qv_i} = S(q) \cap S(v_i)$, respectively. Thus F_{v_i} is a “digonal” facet of $\mathcal{B}(V)$.

The notion of “edge” will be defined in the next paragraph, and Example 5.1 will be generalized in Example 6.2 below.

§ 6. The face structure of Ball polytopes – continuation

Vertices:

Next we turn our attention to the *vertices* of $\mathcal{B}(V)$. As in the previous paragraph we assume (as usual) that $V \subset \mathbb{R}^3$ is finite, non-empty, tight and satisfies $\text{cr}(V) < 1$.

Definition 6.1 (a vertex of $\mathcal{B}(V)$): A boundary point z of $\mathcal{B}(V)$ is a *vertex* of $\mathcal{B}(V)$ if either z belongs to three or more distinct facets of $\mathcal{B}(V)$, in which case z is a *principal vertex*, or $z \in V \cap \mathcal{B}(V)$ and z belongs to exactly two facets of $\mathcal{B}(V)$, in which case z is a *dangling vertex*. Denote by $\text{vert}\mathcal{B}(V)$ the set of vertices of $\mathcal{B}(V)$. In other words, $z \in \text{vert}\mathcal{B}(V)$ if and only if $z \in \mathcal{B}(V)$ and $\|z - p\| = 1$ holds for at least three points $p \in V$, or if $z \in V \cap \mathcal{B}(V)$ and $\|z - p\| = 1$ holds for exactly two points $p \in V$ ¹⁵.

The number of vertices is finite: V is finite, and three distinct unit spheres in $\mathbb{R}^3$ have at most two points in common. Note that if $\#V = 2$, then $\mathcal{B}(V)$ is “lens” shaped with two small caps as facets and no vertices at all. Put $\text{vert}F_p =_{\text{def}} (\text{vert}\mathcal{B}(V)) \cap F_p$ ($p \in V$) and call a point of $\text{vert}F_p$ *vertex* of F_p .

Proposition 6.1: *If $\#V \geq 3$, then the following properties hold true.*

- (i) *A facet F_p ($p \in V$) of $\mathcal{B}(V)$ has at least two vertices which are principal vertices of $\mathcal{B}(V)$.*
- (ii) $\text{vert}F_p \subset \text{relbd}F_p$.

Proof: (i) F_p is strictly spherically convex with a non-empty interior relative to S_p ($=$ the unit sphere with center p). Hence its relative boundary on S_p is a union of at least two closed circular arcs (cf. Theorem 5.4, Proposition 5.1 (ii), and Example 5.1 above), each arc having two endpoints. Each such endpoint is a principal vertex, by Proposition 5.1 (iii).

(ii) Since $\text{relint}F_p = \{x \in S_p : \|x - q\| < 1 \text{ for } q \in V \setminus \{p\}\}$, clearly $\text{relint}F_p \cap \text{vert}\mathcal{B}(V) = \emptyset$, i.e., $F_p \cap \text{vert}\mathcal{B}(V) \subset \text{relbd}F_p$. ■

A facet may contain exactly two principal vertices. In this case and if it contains no dangling vertices, it is a *digonal facet*. Its two “edges” are circular

¹⁵As will become apparent below, a dangling vertex z ($\in V$) always belongs to the relative interior of the intersection $F_p \cap F_q$ of the two facets F_p and F_q , *unlike* the case of 3-polytopes, where every vertex is a “corner” of the body. Nevertheless, this apparently strange notion of “dangling vertex” proves to be fruitful. E.g., in the duality theory developed in § 8 below for every extremal V , $\mathcal{B}(V)$ is self-dual without any artificial restrictions as given in [3], p. 260, lines 19-22 (a restriction that $\mathcal{B}(V)$ has no digonal facets). A similar notion of dangling vertex can also relieve the Bezdek-Naszódi duality theorem in [3], p. 256, Theorem 0.1 (2), from such an artificial restriction (no digonal facets). See footnote 22 below.

arcs (not necessarily of the same radii). In fact, $\mathcal{B}(V)$ may have only two vertices p, q and many digonal facets.

Example 6.1: Let $p = p(0, 0, h)$, $q = q(0, 0, -h)$, $0 < h < 1$, and let $C = C_{pq} = S(p) \cap S(q)$ be a circle in the (x, y) -plane centered at the origin of radius r_h with $0 < r_h = \sqrt{1 - h^2} < 1$. Let $V = \{p_1, p_2, \dots, p_n\}$, $n \geq 3$, be the set of vertices of a regular n -gon inscribed to C . Then V is tight and p, q are principal vertices of $\mathcal{B}(V)$ (there are no other principal vertices), and all n facets $F_{p_1}, F_{p_2}, \dots, F_{p_n}$ of $\mathcal{B}(V)$ are digonal (one can say that $\mathcal{B}(V)$ is “rugby-ball” shaped). When $n \geq 3$ is odd, h with $0 < h < 1$ can be “tuned” so that all the principal diagonals of the regular polygon $[p_1, p_2, \dots, p_n]$ are of length 1, and then all points $p_1, p_2, \dots, p_n$ become dangling vertices of $\mathcal{B}(V)$. (Simple calculation shows that this happens for $h = \sqrt{1 - \frac{1}{4 \cos^2 \frac{180^\circ}{2n}}}$.) In this case the intersection of $\mathcal{B}(V)$ with the (x, y) -plane is a Reuleaux polygon of order n , whose vertices are the dangling vertices of $\mathcal{B}(V)$.

Edges:

Next we turn our attention to the *edges* of $\mathcal{B}(V)$. We would like to define an edge as the intersection $F_p \cap F_q$ of two facets F_p and F_q , provided the intersection is one-dimensional and connected. But it may happen that the intersection $F_p \cap F_q$ is not connected, and that some connectivity components of $F_p \cap F_q$ are singletons (i.e., 0-dimensional); see Example 5.1 above for such a possibility. Hence it is tempting to define an edge of $\mathcal{B}(V)$ as a one-dimensional connectivity component (in the topological sense) of the intersection $F_p \cap F_q$ of two facets. Again this may fail, since a connectivity component of $F_p \cap F_q$ may contain dangling vertices in its relative interior, which is un-acceptable for an edge, of course. This leads us to the following refined definition of an edge.

Definition 6.2 (edge of $\mathcal{B}(V)$): An *edge* of $\mathcal{B}(V)$ is the closure of a connectivity component of $(F_p \cap F_q) \setminus (\text{vert} \mathcal{B}(V))$, where $\{p, q\}$ ranges over all pairs of distinct points of V .

Note that this definition obliterates the possibility of an edge being a singleton.

The intersection $F_p \cap F_q$ satisfies

$$F_p \cap F_q = C_{pq} \cap \bigcap_{v \in V \setminus \{p, q\}} B(v), \quad (9)$$

where C_{pq} is the circle of radius r_{pq} , $0 < r_{pq} = \sqrt{1 - \frac{1}{4}\|p - q\|^2} < 1$, centered in $\frac{1}{2}(p + q)$ lying on the plane that bisects the segments $[p, q]$ perpendicularly. For $\#V = 2$ (i.e., $V = \{p, q\}$), C_{pq} is the only edge of $\mathcal{B}(V)$ (and $\text{vert}\mathcal{B}(V) = \emptyset$). For $\#V \geq 3$, an edge is a relatively closed (short or long) arc of C_{pq} with two endpoints, both being vertices of $\mathcal{B}(V)$; call such a point *vertex* of the edge. (By Proposition 5.1 (i) an edge cannot be the whole of C_{pq} , and by our definition it cannot be a singleton; so an edge has two vertices.) From this we infer:

- (A) $F_p \cap F_q$ is the union of at most $\#V - 2$ edges.
- (B) Every relative boundary point of $F_p \cap F_q$ (relative to C_{pq}) is a principal vertex of $\mathcal{B}(V)$. In particular, an isolated point of $F_p \cap F_q$ (relative to C_{pq}) is a principal vertex. (Such a point is possible only for $\#V \geq 4$, by Proposition 5.1 (iv).)
- (C) A dangling vertex of $\mathcal{B}(V)$ is contained in the relative interior of a non-singleton connectivity component of $F_p \cap F_q$ for some $p, q \in V$.

Proof of (C): A dangling vertex v is a point of V satisfying $\|v - p\| = \|v - q\| = 1$ for some $p, q \in V \setminus \{v\}$, $p \neq q$, and $\|v - w\| < 1$ for $w \in V \setminus \{v, p, q\}$. Thus $v \in F_p \cap F_q = C_{pq} \cap \mathcal{B}(V)$, and a slight movement of v on C_{pq} “back and forth” leaves it in $F_p \cap F_q$, i.e., there is a neighborhood of v relative to C_{pq} contained in $F_p \cap F_q$, i.e., $v \in \text{relint}(F_p \cap F_q)$. ■

- (D) A dangling vertex v of $\mathcal{B}(V)$ is incident with exactly two edges of $\mathcal{B}(V)$, both contained in the connectivity component of v in $C_{pq} \cap \mathcal{B}(V)$.
- (E) For $\#V \geq 3$ the boundary of a facet F_p is a circuit of (at least two) edges separated by vertices. These vertices include at least two principal vertices and possibly dangling vertices appearing in the relative interiors of connectivity components of $F_p \cap F_q$, where q ranges over $V \setminus \{p\}$.

Definition 6.3 (1-skeleton): The geometric graph whose set of vertices is $\text{vert}\mathcal{B}(V)$ and whose edges are the edges of $\mathcal{B}(V)$ is the 1-skeleton of $\mathcal{B}(V)$, denoted by $\text{skel}_1\mathcal{B}(V)$.

As we already saw (cf. Examples 5.1 and 6.1 above), $\text{skel}_1\mathcal{B}(V)$ may have digons, as well as edges of arbitrary (finite) multiplicity. See also Example 6.2 below. For $p \in V$ we have

$$\text{skel}_1\mathcal{B}(V) = (\text{skel}_1\mathcal{B}(V \setminus \{p\}) \cap B(p)) \cup (\text{relbd}F_p), \quad (10)$$

and the union is disjoint except for the set of principal vertices of F_p .

In fact, the set of (at least two) principal vertices of F_p is $(\text{skel}_1\mathcal{B}(V \setminus \{p\})) \cap (\text{relbd}F_p)$ and, by Corollary 6.1 below, for $x \in (\text{vert}\mathcal{B}(V \setminus \{p\})) \cap F_p$ the valence of x in $\text{skel}_1\mathcal{B}(V)$ is greater by 1 than its valence in $\mathcal{B}(V \setminus \{p\})$. Similarly, if $x \in \text{vert}F_p \subset \text{vert}\mathcal{B}(V)$ is of valence ≥ 4 in $\text{skel}_1\mathcal{B}(V)$, then $x \in \text{vert}\mathcal{B}(V \setminus \{p\})$ and x is a principal vertex of $\mathcal{B}(V \setminus \{p\})$.

If $x \in \text{vert}F_p$ and the valence of x in $\text{skel}_1\mathcal{B}(V)$ is 3, then either x lies in the relative interior $\text{relint}e$ of some edge e of $\text{skel}_1\mathcal{B}(V \setminus \{p\})$, or x is a dangling vertex of $\mathcal{B}(V \setminus \{p\})$.

Definition 6.4 (The face complex $\mathcal{SF}(\mathcal{B}(V))$):

The set of faces of $\mathcal{B}(V)$, including facets, edges, vertices, and the improper faces $\mathcal{B}(V)$ and $\emptyset$, is the *spherical face complex* of $\mathcal{B}(V)$ denoted by $\mathcal{SF}(\mathcal{B}(V))$.

The complex $\mathcal{SF}(\mathcal{B}(V))$ is a poset (under inclusion) which may differ considerably from the face lattice of ordinary convex 3-polytopes. The intersection of two facets may have more than one component (cf. Example 5.1 above), there may be digonal facets and there may be only two vertices (cf. Example 6.1 above). Consequently, the set of faces $\mathcal{SF}(\mathcal{B}(V))$, partially ordered by inclusion, need not be a lattice (cf. footnote 5 above), and its 1-skeleton $\text{skel}_1\mathcal{B}(V)$ is a planar graph which is not necessarily 3-connected. Nevertheless, we have

Theorem 6.1: *For a finite set $V \subset \mathbb{R}^3$, $\#V \geq 3$, which is tight with $\text{cr}(V) < 1$, the 1-skeleton $\text{skel}_1(\mathcal{B}(V))$ of $\mathcal{SF}(\mathcal{B}(V))$ is 2-connected (in the sense of*

graph theory)¹⁶.

Proof: We use induction on $n =_{\text{def}} \#V$. For $n = 3$, $\text{skel}_1\mathcal{B}(V)$ has two (principal) vertices (and an edge of multiplicity 3 between them, i.e., 3 edges), which is a 2-connected graph (in the void sense).

Induction step $n \rightarrow n + 1$ ($n \geq 3$): Let $\{x, y, z\} \subset \text{vert}\mathcal{B}(V)$, $x \neq y \neq z \neq x$. It is necessary (and sufficient) to show that there is a path $\gamma_{xy} \subset \text{skel}_1\mathcal{B}(V)$ with endpoints x, y which avoids z .

The basic idea is easily illustrated for the case that $\{x, y, z\} \subset \text{vert}\mathcal{B}(V \setminus \{p\})$ for some $p \in V$. Then there is a path $\tilde{\gamma}_{xy} = \text{skel}_1\mathcal{B}(V \setminus \{p\})$ with endpoints x, y which avoids z (induction hypothesis). If $\tilde{\gamma}_{xy} \subset \text{skel}_1\mathcal{B}(V)$, define $\gamma_{xy} =_{\text{def}} \tilde{\gamma}_{xy}$. Otherwise, we shall modify $\tilde{\gamma}_{xy}$ using (10) (and the discussion thereof). Assuming that $\tilde{\gamma}_{xy}$ is parameterized by t , $a \leq t \leq b$ (“time”), with $\tilde{\gamma}_{xy}(a) = x$ and $\tilde{\gamma}_{xy}(b) = y$, define

$$t_{\min} = \min\{a \leq t \leq b : \tilde{\gamma}(t) \in \text{relbd}F_p\}, \quad x' =_{\text{def}} \tilde{\gamma}_{xy}(t_{\min})$$

and

$$t_{\max} = \max\{a \leq t \leq b : \tilde{\gamma}_{xy}(t) \in \text{relbd}F_p\}, \quad y' =_{\text{def}} \tilde{\gamma}_{xy}(t_{\max}).$$

We say that $x' [y']$ is the *first* [*last*] point of $\tilde{\gamma}_{xy}$ which lies on $\text{relbd}F_p$ (equivalently: on S_p), or that $\tilde{\gamma}_{xy}$ leaves [returns to] $B(p)$ at $x' [y']$ for the first [last] time (possibly $x' = y'$). Since $x', y' \subset (\text{skel}_1\mathcal{B}(V \setminus \{p\})) \cap (\text{relbd}F_p)$, x' and y' are (principal) vertices of $\mathcal{B}(V)$, and they divide $\text{relbd}F_p \subset \text{skel}_1\mathcal{B}(V)$, cf. (10) above) into two paths $A_{x'y'}^1, A_{x'y'}^2$ both with endpoints x', y' ($A_{x'y'}^1 \cap A_{x'y'}^2 = \{x', y'\}$), at least one of which, say $A_{x'y'}^1$, avoids z (if $z \notin \text{relbd}F_p$, then both $A_{x'y'}^i$ ($i = 1, 2$) avoid z). Denote by $\tilde{\gamma}_{xx'} [\tilde{\gamma}_{yy'}]$ the part of $\tilde{\gamma}_{xy}$ between x and x' , $a \leq t \leq t_{\min}$ [y and y' , $t_{\max} \leq t \leq b$] and define $\gamma_{xy} =_{\text{def}} \tilde{\gamma}_{xx'} \circ A_{x'y'}^1 \circ \tilde{\gamma}_{y'y}$

¹⁶In [2], p. 220, Claim 9.4, it is proved that the 1-skeleton of a so-called “standard” ball polytope is 3-connected, where “standard” is a ball polytope every supporting sphere of which (i.e., a sphere which bounds the polytope but does not meet its interior) intersects it in a connected set, i.e., a topological disk (of some dimension, cf. Definition 6.4 there). This assumption is too restrictive since, as Example 5.1 above shows, the 1-skeleton of a ball polytope is *not* 3-connected generically. So the word “standard” there should not be taken literally.

(“ $\circ$ ” means concatenation). Note that we used the facts that $\text{relbd}F_p$ is a circuit in $\text{skel}_1\mathcal{B}(V)$ and that the intersection of a path in $\text{skel}_1\mathcal{B}(V \setminus \{p\})$ with S_p is a vertex of F_p (contained in $\text{relbd}F_p$).

This idea will reappear now in variants adapted to the various cases to be distinguished in the following.

Consider first the case that $\{x, y\} \subset \text{relbd}F_p (\subset \text{skel}_1\mathcal{B}(V))$ for some facet F_p ($p \in V$) of $\mathcal{B}(V)$. Then x, y divide $\text{relbd}F_p$ into two paths A_{xy}^1, A_{xy}^2 both with endpoints x, y ($A_{xy}^1 \cap A_{xy}^2 = \{x, y\}$), at least one of which, say A_{xy}^1 , avoids z (if $z \notin \text{relbd}F_p$, i.e., $z \notin S_p$, then both $A_{xy}^i, i = 1, 2$, avoid z), and define $\gamma_{xy} =_{\text{def}} A_{xy}^1$.

From now on we assume that $\{x, y\} \not\subset F_p$ for all $p \in V$; in particular, x and y are not both vertices of the same edge of $\text{skel}_1\mathcal{B}(V)$ nor do they belong to the same edge of $\text{skel}_1\mathcal{B}(V \setminus \{p\})$. Fix $p \in V$. Our case analysis begins with

Case 1: $\{x, y\} \subset \text{vert}\mathcal{B}(V \setminus \{p\})$.

Subcase 1.1: $z \in \text{vert}\mathcal{B}(V \setminus \{p\})$. This case was fully discussed above.

Subcase 1.2: $z \notin \text{vert}\mathcal{B}(V \setminus \{p\})$. Then $z \in \text{relbd}F_p$, and by (10) (and the discussion thereof) there is an edge $e_{uv} \in \text{skel}_1\mathcal{B}(V \setminus \{p\})$ with endpoints $u, v \in \text{vert}\mathcal{B}(V \setminus \{p\})$ such that $z \in \text{relint}e_{uv}$. Since $\{x, y\} \neq \{u, v\}$ (otherwise x, y lie on the same edge e_{uv} of $\text{skel}_1\mathcal{B}(V \setminus \{p\})$, contrary to our assumption), $u \notin \{x, y\}$ or $v \notin \{x, y\}$, say $v \notin \{x, y\}$. We have now $v \neq x \neq y \neq v$, $\{x, y, v\} \subset \text{vert}\mathcal{B}(V \setminus \{p\})$, and by the induction hypothesis there is a path $\tilde{\gamma}_{xy} \subset \text{skel}_1\mathcal{B}(V \setminus \{p\})$ with endpoints x, y such that $v \notin \tilde{\gamma}_{xy}$. Clearly $z \notin \tilde{\gamma}_{xy}$. If $\tilde{\gamma}_{xy} \cap \text{relbd}F_p = \emptyset$, then $\tilde{\gamma}_{xy} \subset \text{skel}_1\mathcal{B}(V)$, and we define $\gamma_{xy} =_{\text{def}} \tilde{\gamma}_{xy}$.

Assume $\tilde{\gamma}_{xy} \cap \text{relbd}F_p \neq \emptyset$, let $x' [y']$ be the first [last] point of $\tilde{\gamma}_{xy}$ which lies on $\text{relbd}F_p$ (possibly $x' = y'$; surely $z \notin \{x', y'\}$ (clear)), and denote by $\tilde{\gamma}_{xx'} [\tilde{\gamma}_{yy'}]$ the part of $\tilde{\gamma}_{xy}$ between x and x' [y and y']. Points x', y' divide $\text{relbd}F_p$ into two paths $A_{x'y'}^1$ and $A_{x'y'}^2$, both with endpoints x', y' ($A_{x'y'}^1 \cap A_{x'y'}^2 = \{x', y'\}$), at least one of which, say $A_{x'y'}^1$, avoids z . Define $\gamma_{xy} = \tilde{\gamma}_{xx'} \circ A_{x'y'}^1 \circ \tilde{\gamma}_{y'y}$.

Case 2: $\{x, y\} \not\subset \text{vert}\mathcal{B}(V \setminus \{p\})$, say $x \notin \mathcal{B}(V \setminus \{p\})$ (note that the roles of x, y are interchangeable). By (10) (and the discussion thereof) x is a (principal) 3-valent vertex of $\text{skel}_1\mathcal{B}(V)$ lying on $\text{relbd}F_p \subset \text{skel}_1\mathcal{B}(V)$ and $x \in \text{relint}e_{uv}$, where $e_{uv} \in \text{skel}_1(\mathcal{B} \setminus \{p\})$ is an edge not lying on $\text{relbd}F_p$, with

vertices $u, v \in \text{vert}\mathcal{B}(V \setminus \{p\})$. Denote by $e_{xu}[e_{xv}]$ the part (circular arc) of e_{uv} between x and u [x and v]. Since $e_{uv} \not\subset F_p$ is a circular arc, it may have at most one point of intersection with $\text{relbd}F_p$ in addition to x . Call this point w – if it exists – and assume, w.l.o.g., that $w \in e_{xv}$, i.e., $w \notin e_{xu}$ (if w exists; possibly $w = v$). It follows from this convention that $u \notin \text{relbd}F_p$, hence $u \notin S_p$, i.e., either $u \in \text{int}B(p)$ or $u \in \mathbb{R}^3 \setminus B(p)$ and e_{xu} is an edge of $\mathcal{B}(V)$. We also make the following convention and notation: since $x, w \in \text{relbd}F_p$ (if w exists), the points x, w divide $\text{relbd}F_p$ into two paths A_{xw}^1, A_{xw}^2 both with endpoints x, w ($A_{xw}^1 \cap A_{xw}^2 = \{x, w\}$) at least one of which, say A_{xw}^1 , does not contain z in its relative interior (possibly $w = z$; if $z \notin \text{relbd}F_p$, then both A_{xw}^i ($i = 1, 2$) do not contain z). Note that $A_{xy}^1 \subset \text{skel}_1\mathcal{B}(V)$.

Since $y \notin \text{relbd}F_p$ (by our basic assumption x, y do not belong to the same facet of $\mathcal{B}(V)$), $y \in \text{int}\mathcal{B}(p)$ and $y \in \text{vert}\mathcal{B}(V \setminus \{p\})$. If $u = y$, then $e_{xu} = e_{xy}$ is an edge of $\text{skel}_1\mathcal{B}(V)$, a contradiction to our assumption; hence $u \neq y$. If $v = y$, then $e_{xv} = e_{xy}$ is an edge of $\text{skel}_1\mathcal{B}(V \setminus \{p\})$, contradicting our assumption that x, y do not lie on the same edge of $\text{skel}_1\mathcal{B}(V \setminus \{p\})$. To sum up,

$$y \notin \{u, v\}. \quad (11)$$

We distinguish between two subcases.

Subcase 2.1: $z \in \text{vert}\mathcal{B}(V \setminus \{p\})$.

Again we distinguish two subcases.

Sub-subcase 2.1.1: $z \neq u$.

We have $u \neq z \neq y \neq u$ (see (11)) and $\{u, y, z\} \subset \text{skel}_1\mathcal{B}(V \setminus \{p\})$. By the induction hypothesis there is a path $\tilde{\gamma}_{uy} \subset \text{skel}_1\mathcal{B}(V \setminus \{p\})$ with endpoints u, y such that $z \notin \tilde{\gamma}_{uy}$. If $\tilde{\gamma}_{uy} \cap \text{relbd}F_p = \emptyset$ (i.e., $\tilde{\gamma}_{uy} \cap S_p = \emptyset$), then $u \in \text{int}B(p)$ (recall that $y \in \text{int}B(p)$), hence e_{xu} is an edge of $\mathcal{B}(V)$ and we define $\gamma_{xy} =_{\text{def}} e_{xu} \circ \tilde{\gamma}_{uy}$. Assume now that $\tilde{\gamma}_{uy} \cap S_p \neq \emptyset$; call y' the last point of $\tilde{\gamma}_{uy}$ which lies on $\text{relbd}F_p$ (possibly $y' = x$). Since $z \neq x$ and $z \neq y'$, the points x, y' divide the circuit $\text{relbd}F_p$ into two paths $A_{xy'}^1, A_{xy'}^2$ ($A_{xy'}^1 \cap A_{xy'}^2 = \{x, y'\}$) at least one of which, say $A_{xy'}^1$, avoids z (if $z \notin \text{relbd}F_p$, i.e., $z \notin S_p$), then both paths $A_{xy'}^i$ ($i = 1, 2$) avoid z). Denote by $\tilde{\gamma}_{yy'}$ the part of $\tilde{\gamma}_{uy}$ between y' and y and define $\gamma_{xy} =_{\text{def}} A_{xy'}^1 \circ \tilde{\gamma}_{yy'}$.

Sub-subcase 2.1.2: $z = u$.

Then $z \in (\text{relint} B(p)) \cap (\text{vert} \mathcal{B}(V \setminus \{p\}))$. If $v = y$, then $v \in B(p)$ and w is defined; denote by e_{wv} the part of e_{uv} between w and v (in case $w = v$, e_{wv} is just a point); note that e_{wv} is an edge of $\text{skel}_1 \mathcal{B}(V)$ (or a vertex in case $w = v$). Define $\gamma_{xy} = A_{xw}^1 \circ e_{wv}$. Assume now that $v \neq y$. Then $v \neq y \neq z \neq v$ (recall that $z = u \neq v$) and $\{v, y, z\} \subset \text{vert} \mathcal{B}(V \setminus \{p\})$. By the induction hypothesis there is a path $\tilde{\gamma}_{vy} \subset \text{skel}_1 \mathcal{B}(V \setminus \{p\})$ with endpoints v, y which avoids z . If $\tilde{\gamma}_{vy} \cap (\text{relbd} F_p) = \emptyset$, then $v \in B(p)$, w is defined and we define $\gamma_{xy} =_{\text{def}} A_{xw}^1 \circ e_{wv} \circ \tilde{\gamma}_{vy}$.

Assume now that $\tilde{\gamma}_{vy} \cap (\text{relbd} F_p) \neq \emptyset$; call y' the last point of $\tilde{\gamma}_{vy}$ which lies on $\text{relbd} F_p$. Points x, y' divide $\text{relbd} F_p$ into two paths $A_{xy'}^1, A_{xy'}^2$, both with endpoints x, y' ($A_{xy'}^1 \cap A_{xy'}^2 = \{x, y'\}$) both of which avoid z (since $z = u \notin \text{relbd} F_p$). Denote by $\tilde{\gamma}_{y'y}$ the part of $\tilde{\gamma}_{vy}$ between y' and y and define $\gamma_{xy} =_{\text{def}} A_{xy'}^1 \circ \tilde{\gamma}_{y'y}$.

Subcase 2.2: $z \notin \text{vert} \mathcal{B}(V \setminus \{p\})$.

Then $z \in \text{relbd} F_p$, z is a (principal) vertex of valence 3 in $\text{skel}_1 \mathcal{B}(V)$ and there is an edge $e_{st} \in \text{skel}_1 \mathcal{B}(V \setminus \{p\})$ with endpoints $s, t \in \text{vert} \mathcal{B}(V \setminus \{p\})$ such that $z \in \text{relint} e_{st}$ ($z \notin \{s, t\}$). Now we distinguish again two cases.

Sub-subcase 2.2.1: $u \notin \{s, t\}$.

Since $y \neq s$ or $y \neq t$ (or both), we assume, w.l.o.g., that $y \neq s$. Then $s \neq u \neq y \neq s$ (see (11)), $\{u, s, y\} \subset \text{vert} \mathcal{B}(V \setminus \{p\})$, and by the induction hypothesis there is a path $\tilde{\gamma}_{uy} \subset \text{skel}_1 \mathcal{B}(V \setminus \{p\})$ with endpoints u and y which avoids s . Since $e_{st} \notin \tilde{\gamma}_{uy}$ and $\tilde{\gamma}_{uy}$ does not use any edge of $\text{relbd} F_p$, $z \notin \tilde{\gamma}_{uy}$. If $\tilde{\gamma}_{uy} \cap \text{relbd} F_p = \emptyset$, define $\gamma_{xy} =_{\text{def}} e_{xu} \circ \tilde{\gamma}_{uy} \subset \text{skel}_1 \mathcal{B}(V)$.

Assume now that $\tilde{\gamma}_{uy} \cap \text{relbd} F_p \neq \emptyset$ and call y' the last point of $\tilde{\gamma}_{uy}$ which lies on $\text{relbd} F_p$ (equivalently: on S_p). Points x, y' divide $\text{relbd} F_p$ into two paths $A_{xy'}^1, A_{xy'}^2$, both with endpoints x, y' ($A_{xy'}^1 \cap A_{xy'}^2 = \{x, y'\}$), (exactly) one of which, say $A_{xy'}^1$, avoids z . Denote by $\tilde{\gamma}_{y'y}$ the part of $\tilde{\gamma}_{uy}$ between y' and y and define $\gamma_{xy} =_{\text{def}} A_{xy'}^1 \circ \tilde{\gamma}_{y'y}$.

Sub-subcase 2.2.2: $u \in \{s, t\}$, say $u = s$.

Since $u \neq y$ (cf. (11)), we have $u = s \neq y$. Again we distinguish two subcases.

Sub-sub-subcase 2.2.2.1: $y \neq t$.

Then $(s =) u \neq y \neq t \neq u$ and $\{u, y, t\} \subset \text{skel}_1 \mathcal{B}(V \setminus \{p\})$. By the induction hypothesis there is a path $\tilde{\gamma}_{uy} \subset \text{skel}_1 \mathcal{B}(V \setminus \{p\})$ with endpoints u and y which avoids t . The rest of the argument is exactly as in Sub-subcase 2.2.1 above from the wording “Since $\tilde{\gamma}_{wy}$ does not use $e_{st} \dots$ ” to the end “Define $\gamma_{xy} =_{\text{def}} A_{xy'}^1 \circ \tilde{\gamma}_{y'y}$ ”.

Sub-sub-subcase 2.2.2.2: $y = t$.

Then $v \neq t$ (recall that $v \neq y$; cf. (11)), and we have $(s =) u \neq v \neq y (= t) \neq u (= s)$ and $\{u, v, y\} \subset \text{vert} \mathcal{B}(V \setminus \{p\})$. By the induction hypothesis there is a path $\tilde{\gamma}_{vy} \subset \text{skel}_1 \mathcal{B}(V \setminus \{p\})$ with endpoints v and $y (= t)$ which avoids $u (= s)$. Since $e_{st} \notin \tilde{\gamma}_{vy}$ and $\tilde{\gamma}_{vy}$ does not use any edge of $\text{relbd} F_p$, $z \notin \tilde{\gamma}_{vy}$. If $\tilde{\gamma}_{vy} \cap \text{relbd} F_p = \emptyset$, then $v \in \text{int} B(p)$ (recall that $y \in \text{int} B(p)$), w exists (see above) and, by our conventions definition of w above, $z \notin A_{xw}^1$. Call e_{wv} the part of e_{uv} between w and v ($e_{wv} \in \text{skel}_1 \mathcal{B}(V)$) and define $\gamma_{xy} =_{\text{def}} A_{xw}^1 \circ e_{wv} \circ \tilde{\gamma}_{vy}$.

Assume now that $\tilde{\gamma}_{vy} \cap \text{relbd} F_p \neq \emptyset$. Call y' the last point of $\tilde{\gamma}_{vy}$ which lies on $\text{relbd} F_p$, and denote by $\tilde{\gamma}_{y'y}$ the part of $\tilde{\gamma}_{vy}$ between y' and y . Points x, y' divide $\text{relbd} F_p$ into two paths $A_{xy'}^1, A_{xy'}^2$, both with endpoints x and y' ($A_{xy'}^1 \cap A_{xy'}^2 = \{x, y'\}$), (exactly) one of which, $A_{xy'}^1$, say, avoids z . Define $\gamma_{xy} =_{\text{def}} A_{xy'}^1 \circ \tilde{\gamma}_{y'y}$. ■

Remarks 6.1:

1. An alternative approach to Theorem 6.1, via the dual graph of $\mathcal{B}(V)$, can be given along the following lines. First prove as a lemma that the dual graph of $\mathcal{B}(V)$ is connected. Let F, G be two facets of $\mathcal{B}(V)$, let $x \in \text{relint} F$, $y \in \text{relint} G$, and let H be a plane passing through x and y such that $H \cap \text{vert} \mathcal{B}(V) = \emptyset$. Then H intersects $\text{int} \mathcal{B}(V)$ and $\gamma =_{\text{def}} H \cap (\text{bd} \mathcal{B}(V))$ is a (closed) convex curve avoiding $\text{vert} \mathcal{B}(V)$, and x, y divide γ into two paths $\gamma_{xy}^1, \gamma_{xy}^2$ both with endpoints x, y ($\gamma_{xy}^1 \cap \gamma_{xy}^2 = \{x, y\}$). The facets of $\mathcal{B}(V)$ met by γ_{xy}^1 form a succession of facets from F to G , each two successive facets share an edge. This is a path between F and G in the dual graph of $\mathcal{B}(V)$. Let $\{x, y, z\} \subset \text{vert} \mathcal{B}(V)$ ($x \neq y \neq z \neq x$), let F, G be facets of $\mathcal{B}(V)$ containing x and y , respectively, and let $(F_0 = F, F_1, \dots, F_n = G)$ be a path in the dual graph of $\mathcal{B}(V)$ from

F to G (assured by the lemma described above). The union of the relative boundaries of F_i , $\bigcup_{i=0}^n \text{relbd}F_i$, is clearly a connected subgraph of $\text{skel}_1\mathcal{B}(V)$, hence there is a path $\tilde{\gamma}_{xy}$ between x and y in $\text{skel}_1\mathcal{B}(V)$.

If $z \notin \tilde{\gamma}_{xy}$, then $\tilde{\gamma}_{xy}$ is a path in $\text{skel}_1\mathcal{B}(v)$ which avoids z . If $z \in \tilde{\gamma}_{xy}$, then $z \in \text{relbd}F_i$ for some $0 \leq i \leq n$, and $(\text{relbd}F_i) \cap \tilde{\gamma}_{xy}$ is a path A_{ab}^1 on $\text{relbd}F_i$ with endpoints a, b , say ($z \in A_{ab}^1$). Replace A_{ab}^1 by its complementary path $A_{ab}^2 =_{\text{def}} \text{cl}(\text{relbd}F_i \setminus A_{ab}^1)$. Then $z \notin A_{ab}^2$, and the obtained path γ_{xy} connects x and y in $\text{skel}_1\mathcal{B}(V)$ and avoids z .

2. The dual graph of $\mathcal{B}(V)$ is 2-connected as well (as $\text{skel}_1\mathcal{B}(V)$). For a proof one can use an approach similar to the one given in the previous remark. A shorter proof, using already established results, runs as follows. The Bezdek-Naszódi theorem, described in footnote 22 below, assures that the dual graph of $\mathcal{B}(V)$ is isomorphic to the 1-skeleton of the ball polytope $\mathcal{B}(\text{vert}\mathcal{B}(V))$ (in fact, $\mathcal{B}(V)$ and $\mathcal{B}(\text{vert}\mathcal{B}(V))$ have dual face structures), and by Theorem 6.1 above $\text{skel}_1\mathcal{B}(\text{vert}\mathcal{B}(V))$ is 2-connected.

Question 6.1: *Is every 2-connected planar graph realizable as the 1-skeleton of some Ball polytope $\mathcal{B}(V)$ ¹⁷?*

Since $\text{skel}_1\mathcal{B}(V)$ is a finite connected (even 2-connected, by Theorem 6.1) planar graph embedded on the boundary $\text{bd}\mathcal{B}(V)$ of the convex body $\mathcal{B}(V)$, we have:

Proposition 6.2: *If $V \subset \mathbb{R}^3$ is finite with $\#V \geq 3$ and $\text{cr}(V) < 1$, then the face numbers v (vertices), e (edges) and f (facets) of $\mathcal{SF}(\mathcal{B}(V))$ satisfy the Eulerian relation $v - e + f = 2$ ¹⁸.*

Example 6.2: The following Ball polytope $\mathcal{B}(V)$, generalizing Example 5.1 above, exhibits the various notions of faces (vertices, edges, facets) of

¹⁷See [2], Problem 9.5 (p. 221), for a similar question; note that the condition “with no loops” there is superfluous and does not appear in our formulation.

¹⁸This appears as “the Euler-Poincare theorem” in [3], p. 256, with an alleged proof given in p. 260. Instead of saying that this relation follows merely from the planarity and connectivity of the graph $\text{skel}_1\mathcal{B}(V)$, an argument is given based on the sophisticated notion of CW -decomposition from homology theory, not necessary in this simple case. Another drawback of this proof is that the whole face structure of $\mathcal{B}(V)$ is compressed there into two sentences.

$\mathcal{SF}(\mathcal{B}(V))$ defined above in a non-trivial way. We first briefly introduce the basic notion of 2-tuples in equilateral position, which is fully defined in Definition 8.2 below, in connection with the notion of dual arcs.

Let $a, b, x, y \in \mathbb{R}^3$ be four points satisfying $\|x - a\| = \|x - b\| = \|y - a\| = \|y - b\| = 1$. The couple of 2-tuples $(a, b; x, y)$ are in *equilateral position*. Denote by C_{xy} , resp. C_{ab} , the circle centered at $\frac{1}{2}(x + y)$, resp. $\frac{1}{2}(a + b)$, that passes through a, b , resp. x, y .

In Lemma 8.1 below it is proved in detail that if $c \in C_{xy}$, resp. $z \in C_{ab}$, and if both c and z lie on the relative interiors of the short [resp. both lie in the long] arcs of C_{xy} , resp. C_{ab} , determined by a, b , resp. x, y , then $\|c - z\| > 1$. Otherwise $\|c - z\| \leq 1$, and this is satisfied as an equality iff $c \in \{a, b\}$ or $z \in \{x, y\}$. We use this to generalize Example 5.1 above. With the notation $p = p(0, 0, h), q = q(0, 0, -h), 0 < h < 1, C_{pq} = S(p) \cap S(q), p_0, p_1, \dots, p_{n-1} \subset C_{pq} (n \geq 2)$ and $\text{orarc}(p_{i-1}, p_i)$, as introduced there, define n relatively open arcs $A_i, 1 \leq i \leq n$ (replacing the n points $v_i, 1 \leq i \leq n$, of Example 5.1) in the following way: For $1 \leq i \leq n$, if $\text{orarc}(p_{i-1}, p_i)$ is a short [resp. long] arc of C_{pq} , then A_i is the relatively open short [resp. long] arc A_{pq}^i [resp. $C_{p_{i-1}p_i} \setminus A_{pq}^i$] of the circle $C_{p_{i-1}p_i} =_{\text{def}} S(p_{i-1}) \cap S(p_i)$ whose endpoints are p, q . Note that the v_i 's of Example 5.1 satisfy $v_i \in A_i (1 \leq i \leq n)$ and that since $\text{orarc}(p_{i-1}, p_i)$ is a long arc for at most one $i, 1 \leq i \leq n$, at most one $A_i (1 \leq i \leq n)$ is a long arc. It follows from Lemma 8.1 (as explained above) that for $v \in A_i$ the unit ball $B(v, 1)$ misses the relative interior of $\text{orarc}(p_{i-1}, p_i)$ and includes (in its interior) the relative interior of the complementary arc $\text{orarc}(p_i, p_{i-1})$. Let $W \subset \bigcup_{i=1}^n A_i$ be any finite set ($0 \leq \#W < \infty$) and put $V =_{\text{def}} \{p, q\} \cup W$. Then $\text{cr}(V) < 1$, V is tight and, similar to (8) in § 5 above,

$$\begin{aligned} F_p \cap F_q &= C_{pq} \cap \mathcal{B}(V) = \\ &\bigcup \{ \text{orarc}(p_{i-1}, p_i) : W \cap A_i = \emptyset, 1 \leq i \leq n \} \cup \{ p_i : W \cap A_i \neq \emptyset, \\ &\text{and } W \cap A_{i+1} \neq \emptyset, 1 \leq i \leq n \}. \end{aligned} \quad (12)$$

The set of principal vertices $\text{prin-vert} \mathcal{B}(V)$ of $\mathcal{B}(V)$ is contained in the point set $\{p_1, p_2, \dots, p_n = p_0\}$, and for $1 \leq i \leq n$ the point p_i is a principal vertex of $\mathcal{B}(V) \Leftrightarrow W \cap A_i \neq \emptyset$ or $W \cap A_{i+1} \neq \emptyset$ (or both).

Before describing the dangling vertices of $\mathcal{B}(V)$, we describe its 1-skeleton and its facets. The closure of a connectivity component of $\text{skel}_1(\mathcal{B}(V)) \setminus \text{prin-vert}\mathcal{B}(V)$ is a circular arc with two endpoints in $\text{prin-vert}\mathcal{B}(V)$, and it may contain dangling vertices in its relative interior as well. We call such a connectivity component a *hedge* (to differentiate it from “edge”).

A hedge is the union of consecutive edges separated by dangling vertices. Hence a hedge is an edge iff it does not contain a dangling vertex (in its relative interior), and clearly the body of $\text{skel}_1\mathcal{B}(V)$ is the union of its hedges. Every connectivity component of $F_p \cap F_q$ is either a hedge or a principal vertex, and these connectivity components are cyclically ordered on C_{pq} .

Two consecutive points p_{i-1}, p_i of $\{p_1, \dots, p_n = p_0\}$ belong to two consecutive components of $F_p \cap F_q$ iff $A_i \cap W \neq \emptyset$ (in which case $\text{relint}(\text{orarc}(p_{i-1}, p_i))$ is truncated), and if $A_i \cap W \neq \emptyset$, then $p_{i-1}, p_i \in \text{prin-vert}\mathcal{B}(V)$, and p_{i-1}, p_i are connected in $\text{skel}_1\mathcal{B}(V)$ by $\#(A_i \cap W) + 1$ hedges all of which are disjoint from C_{pq} except for their endpoints p_{i-1}, p_i . These $\#(A_i \cap W) + 1$ hedges are the boundaries of $\#(A_i \cap W)$ facets F_v for $v \in A_i \cap W$. Every such facet is “di-hedged”, since its boundary consists of two such hedges (with common endpoints p_{i-1}, p_i); and besides F_p and F_q these are all the facets of $\mathcal{B}(V)$. Thus the hedges of $\mathcal{B}(V)$ are either included in C_{pq} or have endpoints p_{i-1}, p_i for some $1 \leq i \leq n$ such that $A_i \cap W \neq \emptyset$.

A point $v \in W$ is a dangling vertex of $\mathcal{B}(V)$ iff either (i) $v \in C_{pq} \setminus \text{prin-vert}\mathcal{B}(V)$ (in which case $[v, p]$ and $[v, q]$ are edges of $D(V)$), or (ii) v lies in the relative interior of a hedge with endpoints p_{i-1}, p_i such that $A_i \cap W \neq \emptyset$ for some $1 \leq i \leq n$ (i.e., $\text{orarc}(p_{i-1}, p_i)$ is truncated). Dangling vertices satisfying (i) are easy to construct, but the realization of dangling vertices satisfying (ii) turns out to be a delicate question, which we will not address. The following criterion says when p and/or q are dangling vertices.

Criterion: p , resp. q , is a dangling vertex of $\mathcal{B}(V)$ iff $\|p - q\| = 1$, $\#(W \cap C_{pq}) = 1$, and the (unique) arc A_i ($1 \leq i \leq n$) that contains the point $W \cap C_{pq}$ satisfies: $W \cap A_i$ is contained in the closed half space bounded by $\text{aff}C_{pq}$ which contains p , resp. q . (Thus, if both p, q are dangling vertices, then $\#(W \cap A_i) = \#(W \cap C_{pq}) = 1$.)

This criterion finishes our Example 6.2.

The following is fundamental for all later development.

Proposition 6.3: *Let $V \subset \mathbb{R}^3$ be finite with $\#V \geq 3$ and tight with $\text{cr}(V) < 1$. Let F_p be the facet that corresponds to $p \in V$. Then*

- (i) $\text{cone}_p F_p$ is apexed at p .
- (ii) For an edge e of F_p with vertices r and s , the set $\text{cone}_p(r, s)$ is a facet of $\text{cone}_p(\text{vert} F_p)$, and $\text{relint } e \cap \text{cone}_p(\text{vert} F_p) = \emptyset$.
- (iii) The rays $\text{cone}_p r, r \in \text{vert} F_p$, are the extreme rays of $\text{cone}_p(\text{vert} F_p)$, and their cyclic order on $\text{bd}(\text{cone}_p(\text{vert} F_p))$ is compatible with the cyclic order of the vertices of F_p on $\text{relbd} F_p$.

Proof: (i) By Corollary 5.2, F_p is contained in a small cap of S_p .
(ii) There is a facet F_q ($q \in V \setminus \{p\}$) such that $e \subset F_p \cap F_q$ and F_p is contained in the small cap of $S_p = \{x \in \mathbb{R}^3 : \|x - p\| = 1\}$ bounded by the circle $C_{pq} = \text{def}\{x \in \mathbb{R}^3 : \|x - p\| = \|x - q\| = 1\}$ (of radius $0 < r_{pq} =_{\text{def}} \sqrt{1 - \frac{1}{4}\|p - q\|^2} < 1$). Denote this small cap of S_p by $\text{cap}(p)$. $C_{pq} = \text{relbd cap}(p)$, the cap $\text{cap}(p) \supset F_p$ lies beyond the plane $H_{pq} =_{\text{def}} \text{aff} C_{pq}$ relative to p , and the plane $\text{aff}(p, r, s) \supset \text{cone}_p(r, s)$ intersects S_p in a great circle, the short arc between r and s of which is contained in $\text{cap}(p)$. Denote by $\text{arc}(r, s)$ this short arc on $\text{cap}(p)$. Note that $\text{arc}(r, s)$ and e share r and s as common vertices.

Since $\text{arc}(r, s)$ is a geodesic arc on S_p both of whose endpoints belong to F_p , $\text{arc}(r, s) \subset F_p$ (by the spherical convexity of F_p), and since $e \subset F_p$, $\text{sph-conv}(e \cup \text{arc}(r, s)) \subset F_p$ as well (again by the spherical convexity of F_p). Since F_p is strictly spherically convex, $\text{arc}(r, s) \setminus \{r, s\} \subset \text{relint} F_p$, and $\text{sph-conv}(e \cup \text{arc}(r, s)) \setminus e \subset \text{relint} F_p$. Hence

$$\text{sph-conv}(e \cup \text{arc}(r, s)) \cap \text{vert} F_p = \{r, s\} \quad (13)$$

(cf. Proposition 6.1 (ii)); note that $\text{relint } e \cap \text{vert} F_p = \emptyset$, since e is an edge of F_p). The plane $H =_{\text{def}} \text{aff}(p, r, s)$ divides $\mathbb{R}^3$ into two closed half-spaces, one of which contains e ; denote this half-space by H^+ . By $F_p \subset \text{cap}(p)$ we have $\text{vert} F_p \subset \text{cap}(p)$, hence

$$H^+ \cap \text{vert} F_p \subset H^+ \cap \text{cap}(p). \quad (14)$$

Claim:

$$H^+ \cap \text{cap}(p) = \text{sph-conv}(e \cup \text{arc}(r, s)). \quad (15)$$

Proof: $H^+ \cap \text{cap}(p) = (H^+ \cap S_p) \cap \text{cap}(p)$, which is the intersection of two spherically convex sets, hence $H^+ \cap \text{cap}(p)$ is spherically convex and “ $\supset$ ” in (15) follows. Let $m \in H^+ \cap \text{cap}(p)$ and denote by n the center of $\text{cap}(p)$ on S_p . The great circle of S_p passing through n and m cuts $H^+ \cap \text{cap}(p)$ in a short arc $\text{arc}(x, y)$, one of whose endpoints $x \in \text{arc}(r, s)$ and the other endpoint $y \in e$. Hence $m \in \text{arc}(x, y) \subset \text{sph-conv}(e \cup \text{arc}(r, s))$. This proves “ $\subset$ ” in (15), proving the claim.

It follows from (13), (14), and (15) that $\text{vert}F_p \cap H^+ = \{r, s\}$, hence H supports $\text{vert}F_p$ and $\text{cone}_p(r, s)$ is a facet of $\text{cone}_p(\text{vert}F_p)$

(iii) It follows from (ii) that for each edge e of F_p with vertices r, s , say, the rays $\text{cone}_p r$ and $\text{cone}_p s$ are extreme rays of $\text{cone}_p(\text{vert}F_p)$, and since $\text{relbd}F_p$ is a circuit of edges separated by vertices, the cyclic order of the rays $\text{cone}_p r, r \in \text{vert}F_p$ on $\text{bd } \text{cone}_p(\text{vert}F_p)$ is compatible with the cyclic order of the vertices of F_p on $\text{relbd}F_p$. ■

The following proposition gives a simple criterion for an edge of $\mathcal{B}(V)$ being short or long (as an arc of its loading circle); it is needed, e.g., in the proof of Theorem 8.1 below.

Proposition 6.4: *With the usual assumptions on V (see Propositions 6.2 and 6.3 above), let e be an edge of $\mathcal{B}(V)$ with endpoints a, b , and assume that $e \subset C_{xy}$, where $x, y \in V$. Assume that $\#(\text{vert}F_x) \geq 3$ and put $H =_{\text{def}} \text{aff}(x, a, b)$. The following four conditions are equivalent:*

- (i) e is a short arc of C_{xy} .
- (ii) The edge e and the point y lie in different sides of the plane H .
- (iii) The edge e and the set $\text{vert}F_x \setminus \{a, b\}$ lie in different sides of H .
- (iv) The point y and the set $\text{vert}F_x \setminus \{a, b\}$ lie in the same side of H .


Proof: The facet F_x is contained in the small cap $\text{cap}(x) =_{\text{def}} S(x, 1) \cap B(y, 1)$ on the unit sphere $S(x, 1)$, relbd $\text{cap}(x) = C_{xy}$, and both y and $\text{cap}(x)$ lie beyond $\text{aff}C_{xy}$ relatively to x . Put $y^* =_{\text{def}} \text{cone}_x(y) \cap \text{aff}C_{xy}$ and $y^{**} =_{\text{def}} \text{cone}_x(y) \cap \text{cap}(x)$. y^* is the center of the disc $D_{xy} =_{\text{def}} \text{conv}C_{xy}$, and y, y^*, y^{**} lie in the same open half space bounded by H . The center y^* of D_{xy} lies in the plane $\text{aff}C_{xy}$ beyond the line $\text{aff}(a, b)$ relatively to the short arc of C_{xy} whose endpoints are a and b . And since $\text{aff}(a, b) = H \cap \text{aff}C_{xy}$ (in fact, $[a, b] = H \cap D_{xy}$), y^* lies (in the 3-space) beyond H relatively to this short arc. This shows that (i) $\Leftrightarrow$ (ii). Since the facet $F_x \subset \text{cap}(x)$ is strictly spherically convex on $S(x, 1)$, the geodetic line $\gamma = H \cap \text{cap}(x)$ on $\text{cap}(x)$ is contained in $\text{relint}F_x$ except for its endpoints a and b , and γ separates $\text{relbd}F_x \subset \text{cap}(x)$ into two parts; one is e , and the other part, containing $\text{vert}F_x \setminus \{a, b\}$, lies on $\text{cap}(x)$ in the same side of γ as y^{**} . Thus $\text{vert}F_x \setminus \{a, b\}$ and y^{**} lie in the same half space bounded by H , which contains y and y^* as well. This shows that (ii) $\Leftrightarrow$ (iii) $\Leftrightarrow$ (iv). ■

Lemmata 6.1 and 6.4 below establish a connection between the local face structure of $\mathcal{B}(V)$ near a vertex x and the points of V at distance 1 from x .

Lemma 6.1: *Suppose c is a vertex of $\mathcal{B}(V)$, where $V \subset \mathbb{R}^3$, $3 \leq \#V < \infty$, V is tight and $\text{cr}(V) < 1$. Let $\{v \in V : \|v - c\| = 1\} = \{v_1, v_2, \dots, v_m\}$ ($m \geq 3$ for c principal, and $m = 2$ for c dangling, $v_i \neq v_j$ for $1 \leq i < j \leq m$). Then the polyhedral cone $\text{cone}_c(v_1, \dots, v_m)$ is apexed (at c), and the rays $\text{cone}_c(v_i)$, $i = 1, 2, \dots, m$, are the extreme rays of it¹⁹.*

Proof: Put $V(c) =_{\text{def}} \{v_1, \dots, v_m\}$. First we show that the cone $\text{cone}(V(c))$ is apexed (at c) (cf. Definition 4.2). Put $z =_{\text{def}} \text{cc}(V)$, the circumcenter of V , and let r be its circumradius ($r < 1$, by assumption). For $v \in V(c)$ we have $\|v - c\| = 1$ and $\|v - z\| \leq r < 1$. Hence v lies in the open halfspace bounded by the plane H that bisects the segment $[c, z]$ in its midpoint perpendicularly which does not contain c ; see Figure 5. Hence H strictly separates c and $V(c)$.

¹⁹Note the similarity between Lemma 6.1 and Proposition 6.3 (iii) above. The full understanding of this similarity becomes apparent in the rigorous proof of the Bezdek-Naszódi duality theorem; see footnote 22 below.



– Figure 5 –

Thus $\text{cone}_c(V(c))$ is apexed (at c). Next we show that for $v \in V(c)$ the ray $\text{cone}_c(v)$ is extreme in $\text{cone}_c(V(c))$. Assume, by r.a.a., that $\exists v_0 \in V(c)$ such that $\text{cone}_c(v_0)$ is not an extreme ray of $\text{cone}_c(V(c))$. (Clearly, every extreme ray of $\text{cone}_c(V(c))$ is of the form $\text{cone}_c(v)$ for some $v \in V(c)$.) Then $\text{cone}_c(V(c)) = \text{cone}_c(V(c) \setminus \{v_0\})$ and

$$v_0 \in S(c) \cap \text{cone}_c(V(c) \setminus \{v_0\}), \quad (16)$$

($S(c)$ is the unit sphere centered at c). The set $S(c) \cap \text{cone}_c(V(c) \setminus \{v_0\})$ is spherically convex in $S(c)$ by Proposition 4.1 (ii). The r.a.a. is accomplished by the following

Claim: v_0 is inessential in V , i.e.,

$$\mathcal{B}(V \setminus \{v_0\}) = \mathcal{B}(V). \quad (17)$$

Proof: We prove “ $\subset$ ” in (17) (the containment “ $\supset$ ” is clear, as the operation $S \rightarrow \mathcal{B}(S)$ reverses inclusion). Assume $x \in \mathcal{B}(V \setminus \{v_0\})$. In order to prove that $x \in \mathcal{B}(V)$, we use the equivalence $x \in \mathcal{B}(T) \Leftrightarrow T \subset B(x)$, from which it follows that $x \in \mathcal{B}(V \setminus \{v_0\}) \Leftrightarrow V \setminus \{v_0\} \subset B(x)$, and $x \in \mathcal{B}(V) \Leftrightarrow V \subset B(x)$, where $B(x)$ is the unit ball centered at x . Hence

$$V \setminus \{v_0\} \subset B(x) \text{ and, in particular, } V(c) \setminus \{v_0\} \subset B(x). \quad (18)$$

By the first containment in (18) it remains to show that $v_0 \in B(x)$. The set $B(x)$ is spindle-convex. By Proposition 4.2, $S(c) \cap B(x)$ is spherically convex in $S(c)$, hence by the second containment in (18)

$$\text{sph-conv}(V(c) \setminus \{v_0\}) \subset S(c) \cap B(x), \quad (19)$$

(the left-hand side is the spherical convex hull of $V(c) \setminus \{v_0\}$ in $S(c)$).

By Proposition 4.1 (iii), $\text{sph-conv}(V(c) \setminus \{v_0\}) = S(c) \cap \text{cone}_c(V(c) \setminus \{v_0\})$, hence by (19)

$$S(c) \cap \text{cone}_c(V(c) \setminus \{v_0\}) \subset S(c) \cap B(x). \quad (20)$$

It follows from (16) and (20) that $v_0 \in S(c) \cap B(x) \subset B(x)$, hence $v_0 \in B(x)$; this proves the claim.

As already said, the claim brings us to a contradiction (V is tight). ■


Lemma 6.2 (Spatial arm-lemma): *Let C be a circle in $\mathbb{R}^3$ centered at c , let L be the line passing through c perpendicular to $\text{aff}C$, and let $v \in \mathbb{R}^3 \setminus L$. Define a function $f : C \rightarrow \mathbb{R}$ by $f(p) = \|p - v\|$ for $p \in C$. Then*

- a) *f has (exactly) two extremum points, one minimum $p_{\min}$ and one maximum $p_{\max}$ obtained in the following way: let v' be the orthogonal projection of v on the plane $\text{aff}C$ and denote by $\text{pr}_c(\cdot) : \text{aff}C \rightarrow C$ the central projection on C with center c in the plane $\text{aff}C$. Then $p_{\min} = \text{pr}_c(v')$ and $p_{\max} = c + (c - p_{\min}) = 2c - p_{\min}$ (= the diametral point of $p_{\min}$ on the circle C),*
- b) *$f(p)$ strictly increases as the point p moves from $p_{\min}$ to $p_{\max}$ on a half-circle of C with endpoints $p_{\min}$ and $p_{\max}$. (There are two such halves.)*

Proof: By $(f(p))^2 = \|p - v\|^2 = \|p - v'\|^2 + \|v' - v\|^2$ it is sufficient to prove the lemma for the case where $v = v'$ (i.e., $v \in \text{aff}C$). Denote by r the radius of C and by α the small angle between the vectors $v' - c$ ($= v - c$) and $p - c$ ($0 \leq \alpha \leq 180^\circ$). Then

$$(f(p))^2 = \|p - v'\|^2 = \|p - c\|^2 + \|v' - c\|^2 - 2\|p - c\| \cdot \|v' - c\| \cos \alpha \quad (21)$$

(cosinus theorem).




– Figures 6 and 7 –

As the point p moves on one half of the circle C from $p_{\min}$ ($\alpha = 0^\circ$) to $p_{\max}$ ($\alpha = 180^\circ$), $\cos \alpha$ strictly decreases, and the right-hand side of (21) increases. This proves a) and b). ■

Lemma 6.3: *Let C be a circle in $\mathbb{R}^3$ centered at c , and fix $p_0 \in C$. Let H be the plane in $\mathbb{R}^3$ that passes through p_0 and c perpendicularly to $\text{aff}C$. Let H^+ be an open halfspace bounded by H and put $C^+ =_{\text{def}} C \cap H^+$. Then*


$$\begin{aligned}
 &(\forall v \in H^+)(\exists \varepsilon > 0)(\forall p \in C^+) \\
 &(\|p - p_0\| < \varepsilon \Rightarrow \|p - v\| < \|p_0 - v\|).
 \end{aligned}
 \tag{22}$$



– Figure 8 –

Remark: In Figure 8 the plane H is “horizontal” and H^+ lies “above” it, while in Figure 9 below the plane H is perpendicular to the plane of the figure.

Proof: Let v' be the orthogonal projection of v on $\text{aff}C$. Since $v \in H^+$ and H is orthogonal to $\text{aff}C$, $v' \in \text{aff}C \cap H^+$. By the foregoing Lemma 6.2 the function $f(p) =_{\text{def}} \|p - v\|$, $p \in C$, attains its minimum at $p_{\min} = \text{pr}_c(v')$ where $\text{pr}_c(\cdot) : \text{aff}C \rightarrow C$ is the central projection on C with center c ($c \in \text{aff}C$), and it attains its maximum at $p_{\max} = c + (c - p_{\min})$.



– Figure 9 –

Since $v' \in \text{aff}C \cap H^+$, $p_{\min} \in \text{aff}C \cap H^+$ and $p_{\max} \in \text{aff}C \cap H^-$, where H^- is the open half-space bounded by H opposite to H^+ . Denote by C^*

the half-circle of C with endpoints $p_{\min}$ and $p_{\max}$ which contains p_0 (note that $p_0 \notin \{p_{\min}, p_{\max}\}$ since $v \in H^+$; hence $v \notin H$ and $v' \notin \text{aff}\{p_0, c\}$). As the point p moves on C^* starting at $p_{\min}$, the distance $f(p) = \|p - v\|$ strictly increases (by Lemma 6.2). Hence f , restricted to $C^+ \cap C^*$ (= the short arc on C between $p_{\min}$ and p_0), attains its maximum at p_0 . Thus $(\forall p \in C^+ \cap C^*) (\|p - v\| < \|p_0 - v\|)$. This proves (22) with $\varepsilon =_{\text{def}} \|p_{\min} - p_0\|$. $\blacksquare$

Lemma 6.4: *With the assumptions and notation of Lemma 6.1 above assume that the rays $\text{cone}_c(v_i)$ ($i = 1, 2, \dots, m, 1$) appear in this circular order on the boundary of $\text{cone}_c(v_1, \dots, v_m)$. I.e., for $m \geq 3$ the facets of this cone are $\text{cone}_c(v_i, v_{i+1})$ ($i = 1, \dots, m$) (the indices taken modulo m , i.e., $v_{m+1} = v_1$). (For $m = 2$ the cone $\text{cone}_c(v_1, v_2)$ is 2-dimensional.) Then:*

- a) *The facets of $\mathcal{B}(V)$ that meet at c are F_{v_i} , $i = 1, \dots, m$.*
- b) *These facets are separated by edges e_i with $e_i \subset F_{v_i} \cap F_{v_{i+1}}$, $i = 1, \dots, m$.*
- c) *For $m \geq 3$ and $1 \leq i \leq m$ the edge e_i initiates from c on the circle $C_{v_i v_{i+1}} =_{\text{def}} S(v_i) \cap S(v_{i+1})$ to the half space bounded by $H =_{\text{def}} \text{aff}(c, v_i, v_{i+1})$ which contains the set $\{v_1, v_2, \dots, v_{i-1}, \hat{v}_i, \hat{v}_{i+1}, v_{i+2}, \dots, v_m\}$ (and the tangent to e_i at c is perpendicular to H).*
- d) *For $m = 2$ (i.e., when c is a dangling vertex), there are exactly two edges e_1, e_2 incident with c , both lying in $F_{v_1} \cap F_{v_2} \subset C_{v_1 v_2}$, and each edge e_i ($i = 1, 2$) initiates from c to a different side of the plane $H = \text{aff}(c, v_1, v_2)$, e_1 and e_2 have a common tangent at c which is the tangent to $C_{v_1 v_2}$ at c , and this tangent is perpendicular to H .*

Proof: Put $V(c) =_{\text{def}} (v_1, \dots, v_m)$ and define $\delta =_{\text{def}} 1 - \max\{\|v - c\| : v \in V \setminus V(c)\}$. Then $0 < \delta < 1$ and


$$\forall x \in \mathbb{R}^3 : \|x - c\| < \delta \Rightarrow x \in \text{int } \mathcal{B}(V \setminus V(c)). \quad (23)$$

Fix $i \in \{1, \dots, m\}$, and put $C_i =_{\text{def}} C_{v_i v_{i+1}} = S(v_i) \cap S(v_{i+1})$ ($S(v_i)$ and $S(v_{i+1})$ are the unit spheres centered at v_i and v_{i+1} , respectively). C_i is a circle of radius $\sqrt{1 - \frac{1}{4}\|v_{i+1} - v_i\|^2}$ centered at $c_i =_{\text{def}} \frac{1}{2}(v_i + v_{i+1})$, passing through c . Consider two cases:

Case 1 ($m = 2$): Then $\{v_i, v_{i+1}\} = \{v_1, v_2\} = V(c)$, and $\text{cone}_c V(c)$ is 2-dimensional. Since c is a vertex of $\mathcal{B}(V)$ that belongs to exactly two facets of $\mathcal{B}(V)$, c is a dangling vertex of $\mathcal{B}(V)$, hence $c \in V$.

By (23) there is an arc A of length $> 2\delta$ centered at c on the circle $C_i = C_1$ wholly contained in $\mathcal{B}(V \setminus V(c))$. Hence $F_{v_1} \cap F_{v_2} = C_1 \cap \mathcal{B}(V)$ has a non-degenerate connectivity component e which is a closed arc of C_1 containing c in its relative interior. Since c is a dangling vertex of $\mathcal{B}(V)$, e is the union of the two edges of $\mathcal{B}(V)$ that meet at c , and F_{v_1} and F_{v_2} are the only facets of $\mathcal{B}(V)$ that meet at c .

Case 2 ($m \geq 3$): By Lemma 6.1 the cone $\text{cone}_c(V(c))$ is full dimensional, and the rays $\text{cone}_c(v_i), i = 1, \dots, m$, are extreme rays of it. Fix $i \in \{1, \dots, m\}$. The plane $\text{aff}C_i$ bisects the segment $[v_i, v_{i+1}]$ perpendicularly at its midpoint denoted by c_i , hence it is perpendicular to the plane $H_i =_{\text{def}} \text{aff}(v_i, v_{i+1}, c)$. The intersection of these two planes is the line $\text{aff}(c_i, c)$ (this line is a perpendicular bisector of the segment $[v_i, v_{i+1}]$; see Figure 10).



– Figure 10 –

Since the rays $\text{cone}_c(v_i)$ and $\text{cone}_c(v_{i+1})$ are extreme rays of $\text{cone}_c V(c)$, $\text{cone}_c(v_i, v_{i+1})$ is a (2-dimensional) facet of $\text{cone}_c(V(c))$, hence the plane H_i of this facet strictly supports $V(c) \setminus \{v_i, v_{i+1}\}$.

Denote by H_i^+ , resp. H_i^- , the open halfspace bounded by H_i which contains,

resp. does not contain, $V(c) \setminus \{v_i, v_{i+1}\}$ and put $C_i^+ = C_i \cap H^+$. By Lemma 6.3 (with $C = C_i$, $p_0 = c$ and $v = v_j$)

$$\begin{aligned} & (\forall v_j \in V(c) \setminus \{v_i, v_{i+1}\})(\exists \varepsilon_j > 0)(\forall p \in C_i^+) \\ & ((\|p - c\| < \varepsilon_j) \Rightarrow (\|p - v_j\| < 1)). \end{aligned} \quad (24)$$

Put $\varepsilon' =_{\text{def}} \min\{\varepsilon_j : j \in \{1, \dots, m\} \setminus \{i, i+1\}\}$ and $\varepsilon'' =_{\text{def}} \min\{\varepsilon', \delta\}$. Then (23) and (24) imply

$$(\forall v \in V \setminus \{v_i, v_{i+1}\})(\forall p \in C_i^+)((\|p - c\| < \varepsilon'') \Rightarrow (\|p - v\| < 1)). \quad (25)$$

It follows that there is an arc A of length $> \varepsilon''$ on C_i with one endpoint at c such that

$$\text{relint} A \subset H_i^+ \cap \text{int } \mathcal{B}(V \setminus \{v_i, v_{i+1}\}). \quad (26)$$

Thus $C_i \cap \text{bd} \mathcal{B}(V)$ has a component which contains A , i.e., there is an edge e_i of $\mathcal{B}(V)$ on C_i which is incident with c , and e_i is common to F_{v_i} and $F_{v_{i+1}}$. This proves a), and e_i satisfies all the requirements of b) and c), as it is easy to check. $\blacksquare$

From Lemma 6.4 we have

Corollary 6.1: *If $V \subset \mathbb{R}^2$ with $\#V \geq 3$ is finite, tight, and $\text{cr}(V) < 1$, then the valence of a vertex x of $\mathcal{B}(V)$ in the 1-skeleton of $\mathcal{SF}(\mathcal{B}(V))$ is equal to the number of points of V at distance 1 from x ²⁰.*

§ 7. On $\mathcal{B}(V)$ for an extremal set V , and an extended GHS-Theorem

In this paragraph we use the foregoing results to establish an extended form of the GHS-theorem. Recall that an *extremal configuration* (for the Vázsonyi

²⁰This corollary was observed (without a proof) by Grünbaum, Heppes, and Straszewicz independently of each other for the case that V is extremal (for the Vázsonyi problem). The corollary generalizes their observation to any feasible set V (i.e., $3 \leq \#V < \infty$, V is tight and satisfies $\text{cr}(V) < 1$).

problem) is a finite set $V \subset \mathbb{R}^3$ with $\#V = n \geq 4$ such that $e(V) = e(3, n)$. We **ignore** now the GHS-Theorem ($e(3, n) = 2n - 2$) in order to achieve a detailed proof of an extended form of it. As said in the introduction, this is essentially an elaboration of the proof given by Grünbaum, Heppes, and Straszewicz. The following result was derived in Proposition and Definition 2.1 above as a consequence of the GHS-Theorem. Now we derive it independently of it.

Proposition 7.1: *Let $V \subset \mathbb{R}^3$, $\#V = n \geq 4$, be an extremal configuration for the Vázsonyi problem. Then the valence of a point $v \in V$ in the diameter graph $D(V)$ is ≥ 2 , i.e., if $\text{diam}V = 1$, then $\text{dist}(v, V \setminus \{v\}) = 1$, and this distance is attained at least twice.*

Proof: Assume, without loss of generality, that $\text{diam}V = 1$. If there is a point $v \in V$ such that $\text{dist}(v, V \setminus \{v\}) < 1$, then choose a line l through v and move v on l until it reaches for the first time a point v' such that $\text{dist}(v', V \setminus \{v\}) = 1$. Put $V' =_{\text{def}} (V \setminus \{v\}) \cup \{v'\}$. Then $\text{diam}V' = \text{diam}V = 1$, $\#V' = \#V = n$, and $e(V') > e(V)$, contrary to our hypothesis (that V is extremal).

A similar contradiction is obtained when $\text{dist}(v, V \setminus \{v\}) = 1$ and this distance is attained only once, i.e., if there is a point $p \in V$ such that $\|v - p\| = 1$ and $\text{dist}(v, V \setminus \{p, v\}) < 1$. Let q be the symmetric point of p relative to v on the line $\text{aff}(p, v)$, and let C be a semi-circle of radius 1 with center v and endpoints p, q . Since v is the center of the segment $[p, q]$ whose length is 2 and since $\text{dist}(p, V \setminus \{p, v\}) \leq 1$, clearly $\text{dist}(q, V \setminus \{p, v\}) > 1$.

Move p on C until it reaches for the first time a point p' such that $\text{dist}(p', V \setminus \{p, v\}) = 1$. This will happen before p reaches q since $\text{dist}(q, V \setminus \{p, v\}) > 1$. Put $V' =_{\text{def}} (V \setminus \{p\}) \cup \{p'\}$. Then $\text{diam}V' = \text{diam}V = 1$, $\#V' = \#V$, and $e(V') > e(V)$, a contradiction. Thus $\text{dist}(v, V \setminus \{v\}) = 1$, and this distance is achieved at least twice. ■

Proposition 7.2: *Let $V \subset \mathbb{R}^3$ with $\infty > \#V = n \geq 4$ be an extremal configuration for the Vázsonyi problem such that $\text{diam}V = 1$. Then:*

- a) $\mathcal{B}(V)$ is full-dimensional and V is tight.
- b) $\mathcal{B}(V)$ has n facets.

c) $V \subseteq \text{vert}\mathcal{B}(V)$.

Proof:

- (a) The first part of a) follows from Theorem 5.2 a). The second part of a) follows from Theorem 5.2 b) using Proposition 7.1 above.
- (b) This follows from the fact that V is tight.
- (c) The valence of $v \in V$ in $D(V)$ is ≥ 2 for $v \in V$ (Proposition 7.1). If v has valence 2 in $D(V)$, then it belongs to exactly two facets of $\mathcal{B}(V)$, i.e., it is a dangling vertex of $\mathcal{B}(V)$. Otherwise v is of distance 1 from at least three distinct points $p, q, r \in V$, i.e., $v \in F_p \cap F_q \cap F_r$ and v is a principal vertex of $\mathcal{B}(V)$. ■

Proposition 7.3: *Let $V \subset \mathbb{R}^3$ be tight with $\infty > \#V = n \geq 1$, $\text{cr}(V) < 1$, and assume that $V \subseteq \text{vert}\mathcal{B}(V)$. Then $\text{diam } V = 1$,*

$$e(V) \leq 2n - 2, \tag{27}$$

and this inequality is satisfied as an equality iff $V = \text{vert}\mathcal{B}(V)$.

Proof: Since $\text{cr}(V) < 1$, $\mathcal{B}(V)$ is full-dimensional. Since $V \subseteq \text{vert}\mathcal{B}(V)$, $\mathcal{B}(V)$ has a non-empty set of vertices, hence $\#V = n \geq 3$. The face numbers of $\mathcal{SF}(\mathcal{B}(V))$ satisfy the Eulerian relation $v - e + f = 2$ (cf. Proposition 6.2). Here v is the number of (principal or dangling) vertices of $\mathcal{B}(V)$, e is the number of edges, and f is the number of facets which is a-priori $\leq n$, but since V is tight, $f = n$. Since $V \subseteq \text{vert}\mathcal{B}(V) \subset \mathcal{B}(V)$, $\text{diam} V \leq 1$. Each point $v \in V$, being a vertex of $\mathcal{B}(V)$, is at distance 1 from at least two points of V , hence $\text{diam} V = 1$. Put $\tilde{V} =_{\text{def}} \text{vert}\mathcal{B}(V)$. Thus $V \subseteq \tilde{V}$ and $\#\tilde{V} = v = n + m$ for some $m \geq 0$. The number e of edges of $\mathcal{B}(V)$ is one-half the sum of valences of the vertices of $\mathcal{B}(V)$ in the 1-skeleton of $\mathcal{SF}(\mathcal{B}(V))$. The valence of each vertex x of $\mathcal{B}(V)$, henceforth denoted by $\text{val}(x)$, is equal

to the number of points of V at distance 1 from x (cf. Corollary 6.1 in the end of § 6). Thus

$$\begin{aligned}
2e &= \sum \{\text{val}(x) : x \in \tilde{V}\} = \\
&= \sum \{\text{val}(x) : x \in V\} + \sum \{\text{val}(x) : x \in \tilde{V} \setminus V\} \\
&\geq \sum \{\text{val}(x) : x \in V\} + 3 \cdot \#(\tilde{V} \setminus V) \\
&= 2 \cdot \#\{\{u, w\} \subset V : \|u - w\| = 1\} + 3m \\
&= 2 \cdot e(V) + 3m.
\end{aligned}$$

It follows, using the Eulerian relation $v - e + f = 2$, that

$$\begin{aligned}
2 \cdot e(V) &\leq 2e - 3m \\
&= 2(v + f - 2) - 3m \quad (\text{by } v - e + f = 2) \\
&= 2(n + m + f - 2) - 3m \\
&= 2(2n - 2) - m \quad (\text{since } f = n).
\end{aligned}$$

Thus $e(V) \leq 2n - 2$, and this is satisfied as an equality iff $m = 0$, i.e., iff $V = \text{vert}\mathcal{B}(V)$. ■

This leads to

Theorem 7.1 (Extended GHS-Theorem): *Let $V \subset \mathbb{R}^3$ be finite with $\#V = n \geq 4$ and $\text{diam}V = 1$. The following three statements are equivalent:*

- (i) V is extremal for the Vázsonyi problem, i.e., $e(V) = e(3, n)$;
- (ii) $e(V) = 2n - 2$;
- (iii) V is tight and $V = \text{vert}\mathcal{B}(V)$ ²¹.

²¹(a) The equivalence of statements (i) and (ii) is what we call the GHS-Theorem (see Corollary 7.1 below), and their equivalence to (iii) is what “Extended” stands for in the name of the theorem.

(b) Sallee in [25], p. 318, calls a Ball polytope satisfying (iii) “a frame of a Reuleaux polytope”, and he points on their usefulness to construct concrete bodies of constant width in $\mathbb{R}^3$. This is analogous to the planar case, where Reuleaux polygons are paradigmatic for (planar) bodies of constant width. (He does not point on the connection of his “frames of Reuleaux polytopes” with the Vázsonyi problem, however.) We may come to this topic in the mentioned subsequent paper.

Proof: $\text{diam}V = 1 \Rightarrow \text{cr}(V) < 1$ (Theorem 5.2, a)), hence $\mathcal{B}(V)$ is full-dimensional.

Proof of (i) $\Rightarrow$ (ii) and (i) $\Rightarrow$ (iii): Assume (i). By Proposition 7.2 a), c) the set V is tight, and $V \subseteq \text{vert}\mathcal{B}(V)$. Hence, by Proposition 7.3, $e(V) \leq 2n - 2$. On the other hand, $e(V) = e(3, n) \geq 2n - 2$, by the examples given in § 1. Thus $e(V) = 2n - 2$ and, again by the suffix to Proposition 7.3, $V = \text{vert}\mathcal{B}(V)$.

Proof of (ii) $\Rightarrow$ (i): By (i) $\Rightarrow$ (ii) proved above $e(3, n) = 2n - 2$. Hence $e(V) = e(3, n)$, i.e., V is extremal.


Proof of (iii) $\Rightarrow$ (ii): Assume that V is tight and $V = \text{vert}\mathcal{B}(V)$. Thus $V \subseteq \text{vert}\mathcal{B}(V)$ and the conditions of Proposition 7.3 are fulfilled. By the suffix to this proposition $V = \text{vert}\mathcal{B}(V) \Rightarrow e(V) = 2n - 2$. ■

From Theorem 7.1 follows

Corollary 7.1 (GHS-Theorem): $e(3, n) = 2n - 2$.

Corollary 7.2: *Assume that $V \subset \mathbb{R}^3$, $4 \leq \#V < \infty$, $\text{diam}V = 1$ and V is extremal for the Vázsonyi problem. Then every edge of $\mathcal{B}(V)$ is a short arc of its loading circle.*

Proof: Let e be an edge of $\mathcal{B}(V)$ with endpoints $a, b \in \text{vert}\mathcal{B}(V) = V$. There are points $x, y \in V$ such that $e \subset C_{xy} =_{\text{def}} S_x \cap S_y$ (S_x, S_y are the spheres of radius 1 centered at x and y , respectively). C_{xy} is a circle of radius $r_{xy} = \sqrt{1 - \frac{1}{4}\|x - y\|^2}$ centered at $p = \frac{1}{2}(x + y)$, $a, b \in C_{xy}$, and e is either the short arc between a and b on C_{xy} , or the respective long arc. We will show that the second case is impossible.



– Figure 11 –

Denote by c_1 [resp. c_2] the midpoint of the short [resp. long] arc between a and b on C_{xy} (cf. Figure 11). c_1 and c_2 are diametrical points of C_{xy} . Since $\|x - y\| \leq 1$, $r_{xy} = \sqrt{1 - \frac{1}{4}\|x - y\|^2} \geq \frac{\sqrt{3}}{2} > \frac{1.7}{2} = 0.85$; hence $\text{diam} C_{xy} = 2r_{xy} \geq 1.7 > 1$. Since c_2 is the midpoint of the long arc between a and b on C_{xy} , clearly $\angle c_2pa > 90^\circ$. It follows that

$$\begin{aligned} \|c_2 - a\|^2 &= \|c_2 - p\|^2 + \|a - p\|^2 - 2\|c_2 - p\| \cdot \|a - p\| \cdot \cos \angle c_2pa \\ &> \|c_2 - p\|^2 + \|a - p\|^2 = 2(r_{xy})^2 \geq 2 \cdot \left(\frac{\sqrt{3}}{2}\right)^2 = \frac{3}{2} > 1. \end{aligned}$$

Hence $\|c_2 - a\| > 1$ and $c_2 \notin B(a, 1)$. Since $a \in \text{vert} \mathcal{B}(V) = V$ (by Theorem 7.1 (iii)), this implies $c_2 \notin \mathcal{B}(V)$, hence $c_2 \notin e$, i.e., e is *not* the long arc between a and b on C_{xy} . Thus e is the short arc between a and b on C_{xy} . ■

§ 8. The canonical self-duality φ of $\mathcal{SF}(\mathcal{B}(V))$ for an extremal set V

In this paragraph we assume that V is a set of n points in $\mathbb{R}^3$, $4 \leq n < \infty$, $\text{diam} V = 1$, and V is extremal for the Vázsonyi problem. Under these assumptions V coincides with the set of vertices of its Ball polytope $\mathcal{B}(V)$ (Extended GHS-Theorem, Theorem 7.1 (iii) above). The poset $\mathcal{SF}(\mathcal{B}(V))$, partially ordered by inclusion, consists of vertices, edges and facets of $\mathcal{B}(V)$ (and, if one wishes, the improper faces $\emptyset$ and $\mathcal{B}(V)$ as well).

Definition 8.1: An *involutory self-duality* (or *antimorphism*) of $\mathcal{SF}(\mathcal{B}(V))$ is an order reversing map (i.e., a duality) $\varphi : \mathcal{SF}(\mathcal{B}(V)) \longrightarrow \mathcal{SF}(\mathcal{B}(V))$ of order two ($\varphi^2 = \text{identity}$).

The correspondence $\{x\} \xleftrightarrow{\varphi} F_x$ ($x \in V$) (and, if necessary, $\emptyset \xleftrightarrow{\varphi} \mathcal{B}(V)$) is, in fact, order reversing, i.e.,

$$\begin{aligned} \{x\} \subset F_y &\Leftrightarrow \|x - y\| = 1 \quad \Leftrightarrow \quad \{y\} \subset F_x \\ &\Leftrightarrow \quad \varphi(F_y) \subset \varphi(\{x\}). \end{aligned}$$

And it is, of course, of order two. This correspondence can be extended to an involutory self-duality of $\mathcal{SF}(\mathcal{B}(V))$ if and only if the following statement holds.

Theorem 8.1 (dual edges): *Let e be an edge of $\mathcal{B}(V)$, with endpoints a and b , and assume that e is included in the facets F_x and F_y of $\mathcal{B}(V)$ ($x, y \in V, x \neq y$). Then there is another edge e' of $\mathcal{B}(V)$, with endpoints x and y , included in the facets F_a and F_b . The edge e' is, of course, uniquely determined by e .*

This theorem is indeed true. Its proof takes some effort, however, and will be postponed (until after Lemma 8.1 below). From this theorem we conclude

Corollary – Definition 8.1 (canonical self-duality φ of $\mathcal{SF}(\mathcal{B}(V))$):

If $V \subset \mathbb{R}^3$, with $\#V \geq 4$ and $\text{diam}(V) = 1$, is an extremal configuration for the Vázsonyi problem, then the correspondence $x \xleftrightarrow{\varphi} F_x$ ($x \in V$) has a unique edge-extension $\tilde{\varphi}, e \xleftrightarrow{\tilde{\varphi}} e'$, which is an involutory self-duality of $\mathcal{SF}(\mathcal{B}(V))$. This self-duality $\tilde{\varphi}$ is the canonical self-duality of $\mathcal{SF}(\mathcal{B}(V))$. Abusing notation we drop henceforth the tilde $\sim$ and denote $\tilde{\varphi}$ by φ ^{22 23}.

²²This self-duality was partially observed by Sallee in [25], p. 322, Remark 2). Our $\mathcal{B}(V)$ is called there “a frame of a Reuleaux polytope” (cf. footnote 20 above), and something like our Theorem 8.1 above is formulated in (6.2) in p. 320 there. His treatment is sketchy, however, overlooking some delicate points.

²³Also, this corollary turns out to be a partially special case of the following remarkable duality theorem of [3], p. 256: If $\mathcal{B}(V)$ is a ball polytope without digonal facets (in $\mathbb{R}^3$), then $\mathcal{B}(V)$ and $\mathcal{B}(\text{vert}\mathcal{B}(V))$ have dual face structures. The proof given there (pp. 261–262) is un-convincing. To see this, it is sufficient to point out that even such most basic properties as enunciated in Proposition 6.3 (iii) and Lemmata 6.1, 6.3, and 6.4 above are

The proof of Theorem 8.1 makes use of a geometric fact for which we need the following

Definition 8.2 (dual arcs): Assume that a, b, x, y are four distinct points in $\mathbb{R}^3$, such that $\|x - a\| = \|x - b\| = \|y - a\| = \|y - b\| = 1$. The (spatial) quadrilateral (a, x, b, y) is an *equilateral quadrilateral*, and the couple of 2-tuples $(a, b; x, y)$ are in *equilateral position*. Denote by C_{xy} , resp. C_{ab} , the circle centered at $\frac{1}{2}(x + y)$, resp. $\frac{1}{2}(a + b)$, that passes through a and b , resp. x and y . Let A_{ab} [resp. A_{xy}] be the short open circular arc of C_{xy} [resp. C_{ab}] with endpoints a, b [resp. x, y]. The circular arcs A_{ab} and A_{xy} are *short dual arcs*. The complementary arcs $C_{xy} \setminus (\text{relint} A_{ab})$ and $C_{ab} \setminus (\text{relint} A_{xy})$ are *long dual arcs*.

Note that the radii r_{ab} and r_{xy} of the circles C_{ab} and C_{xy} , respectively, are both < 1 and, of course, these radii may be different from each other. Note also that the planes $\text{aff} C_{xy}$ and $\text{aff} C_{ab}$ are perpendicular to each other.

The following lemma partially appears as Lemma 4.2 in [26], with an analytic proof. We extend it here and give a purely geometric proof.


Lemma 8.1 (pointwise distance between dual arcs and their complements): *Let A_{ab} and A_{xy} be two (circular) short dual arcs, lying on the circles C_{xy} and C_{ab} , respectively, and let $c \in C_{xy}$ and $z \in C_{ab}$.*

- a) *If $c \in \text{relint} A_{ab}$ [resp. $c \in C_{xy} \setminus A_{ab}$] and $z \in \text{relint} A_{xy}$, then $\|c - z\| > 1$ [resp. $\|c - z\| < 1$].*
- b) *If $c \in C_{xy} \setminus A_{ab}$ and $z \in C_{ab} \setminus A_{xy}$, then $\|c - z\| > 1^{24}$.*

nowhere mentioned there, without which a rigorous treatment of the face structure of $\mathcal{B}(V)$ is impossible. Also the treatment of the edge-edge duality in p. 262 there is unconvincing merely because of the lack of the important geometric inequality enunciated in Lemma 8.1 below (pointwise distance between dual arcs), which takes care of the delicate possibility that what is called the “desired edge” (p. 262, line 16) contains no point of X ($=$ our V) in its relative interior. Also dual edges of $\mathcal{B}(V)$ and $\mathcal{B}(\text{vert} \mathcal{B}(V))$ are both either short or long (as arcs of their corresponding loading circles) – which follows from Proposition 6.4 and Lemma 6.4 above, a fact not mentioned there. Let us also mention that by defining a variant of our dangling vertices (cf. Definition 6.1 above) one can relieve this Bezdek-Naszódi duality theorem from the manifestly artificial assumption that $\mathcal{B}(V)$ contains no digonal facets.

²⁴This lemma is fundamental for ball polytopes. E.g., it has to be used in any rigorous

In other words: If $c \in C_{xy}$ resp. $z \in C_{ab}$, and if either both c resp. z lie on the relative interiors of the short [resp. long] arcs of C_{xy} resp. C_{ab} , with endpoints a, b resp. x, y , then $\|c - z\| > 1$. Otherwise $\|c - z\| \leq 1$, and this is satisfied as an equality iff $c \in \{a, b\}$ or $z \in \{x, y\}$ (or both).



– Figure 12 –

Proof: The line $\text{aff}(\frac{1}{2}(a+b), \frac{1}{2}(x+y))$ passing through the center $\frac{1}{2}(a+b)$ of the circle C_{ab} and through the midpoint $\frac{1}{2}(x+y)$ of its chord $[x, y]$ intersects the circle C_{ab} in two diametrical points z_1 and z_2 ; we choose the notation so that $z_1 \in A_{xy}$ (in fact, z_1 is the midpoint of the arc A_{xy}) and $z_2 \in C_{ab} \setminus A_{xy}$ (z_2 is the midpoint of this long arc of C_{ab}). The planes $H_{xy} =_{\text{def}} \text{aff} C_{xy}$ and $H_{ab} =_{\text{def}} \text{aff} C_{ab}$ intersect perpendicularly in the line $\text{aff}(z_1, z_2)$, and $\frac{1}{2}(a+b)$ is the midpoint of both segments $[a, b]$ and $[z_1, z_2]$ (cf. Figure 12).

Claim: $\text{aff}(a, b) \perp \text{aff}(z_1, z_2)$

Proof: Since $z_1, z_2 \in \text{aff}\{\frac{1}{2}(x+y), \frac{1}{2}(a+b)\}$, it is sufficient to show that

proof of the fundamental Bezdek-Naszódi duality theorem (cf. the former footnote), especially for the edge-edge duality, alas it is lacked in [3], p. 262, which is one of the reasons we find this proof un-convincing.

$\langle a - b, \frac{1}{2}(x + y) - \frac{1}{2}(a + b) \rangle = 0$. This is equivalent to

$$\begin{aligned}
& \langle a - b, x - a + y - b \rangle = 0 \\
\Leftrightarrow & \langle 2a - 2b, x - a + y - b \rangle = 0 \\
\Leftrightarrow & \langle a - x + a - y + x - b + y - b, x - a + y - b \rangle = 0 \\
\Leftrightarrow & \langle a - x, x - a \rangle + \langle y - b, y - b \rangle + \langle a - x, y - b \rangle + \langle y - b, x - a \rangle \\
& + \langle a - y + x - b, x - a + y - b \rangle = 0 \\
\Leftrightarrow & -1 + 1 + \langle y - b, a - x + x - a \rangle \\
& + \langle a - y + x - b, x - b + y - a \rangle = 0 \\
\Leftrightarrow & \langle a - y + x - b, x - b + y - a \rangle = 0 \\
\Leftrightarrow & \langle a - y, y - a \rangle + \langle x - b, x - b \rangle \\
& + \langle a - y, x - b \rangle + \langle x - b, y - a \rangle = 0 \\
\Leftrightarrow & -1 + 1 + \langle x - b, a - y + y - a \rangle = 0 \\
\Leftrightarrow & \langle x - b, 0 \rangle = 0 \Leftrightarrow 0 = 0,
\end{aligned}$$

which proves the claim.

Let $\text{pr}(\cdot) : \mathbb{R}^3 \rightarrow H_{ab}$ be the orthogonal projection on the plane H_{ab} . Since $H_{ab} \perp H_{xy}$,

$$\text{pr}(C_{xy}) \subset H_{ab} \cap H_{xy} = \text{aff}(z_1, z_2), \quad (28)$$

and by the foregoing claim

$$\text{pr}(a) = \text{pr}(b) = \frac{1}{2}(a + b). \quad (29)$$

Since A_{ab} is a short arc on C_{xy} , $\text{relint} A_{ab}$ lies beyond the line $\text{aff}(a, b)$ relatively to the center $\frac{1}{2}(x + y)$ of C_{xy} in the plane H_{xy} . Hence, by (29),

$$\text{pr}(A_{ab}) \subset -\text{cone}_{\frac{1}{2}(a+b)} \frac{1}{2}(x + y) \quad (30)$$

and

$$\text{pr}(C_{xy} \setminus A_{ab}) \subset \text{cone}_{\frac{1}{2}(a+b)} \frac{1}{2}(x + y). \quad (31)$$

Similarly, since A_{xy} is a short arc on C_{ab} , $\text{relint}A_{xy}$ lies beyond the line $\text{aff}(x, y)$ relatively to the center $\frac{1}{2}(a + b)$ of C_{ab} in the plane H_{ab} . Hence, in particular,

$$z_1 \in \text{cone}_{\frac{1}{2}(a+b)} \frac{1}{2}(x + y) \quad (32)$$

and

$$z_2 \in \text{cone}_{\frac{1}{2}(a+b)} \frac{1}{2}(x + y). \quad (33)$$

It follows from (30), (31), (32), and (33) that

$$\text{pr}(A_{ab}) \subset \text{cone}_{\frac{1}{2}(a+b)} z_2 \quad (34)$$

and

$$\text{pr}(C_{xy} \setminus A_{ab}) \subset \text{cone}_{\frac{1}{2}(a+b)} z_1. \quad (35)$$

Back to the proof of Lemma 8.1, let $c \in \text{relint}A_{ab}$ [resp. $c \in C_{xy} \setminus A_{ab}$]. Then, by (34) and (35), $c' =_{\text{def}} \text{pr}(c)$ lies on $\text{cone}_{\frac{1}{2}(a+b)} z_2$ [resp. $\text{cone}_{\frac{1}{2}(a+b)} z_1$]. Hence, by Lemma 6.2 (Spatial arm-lemma), $f : C_{ab} \rightarrow \mathbb{R}$, defined by $f(z) =_{\text{def}} \|z - c\|$ ($z \in C_{ab}$), attains its minimum at z_2 [resp. z_1] and its maximum at z_1 [resp. z_2], and as the point z moves on half of the circle C_{ab} from z_2 to z_1 , $f(z)$ increases [resp. decreases]. Since for $z = x$ we have $f(x) = \|x - c\| = f(y) = \|y - c\| = 1$ (recall that $c \in C_{xy}$ whether $c \in A_{ab}$ or $c \in C_{xy} \setminus A_{ab}$) as z moves from z_2 to z_1 on half of the circle C_{ab} that contains x we have by increasing [resp. decreasing] monotonicity of $f(z)$ that $f(z) = \|z - c\| < \|x - c\| = 1$ [resp. $f(z) = \|z - c\| > \|x - c\| = 1$] when z lies on the short arc $\text{arc}(z_2, x)$ of C_{ab} , and $f(z) = \|z - c\| > \|x - c\| = 1$ [resp. $f(z) = \|z - c\| < \|x - c\| = 1$] when z lies on the short arc $\text{arc}(x, z_1)$ of C_{ab} . Similar inequalities hold as z moves from z_2 to z_1 on half of the circle C_{ab} that contains y . This proves Lemma 8.1 a).

To prove b), assume that the points c, z are as stated in the premises. Now we have $c' =_{\text{def}} \text{pr}(c) \in \text{cone}_{\frac{1}{2}(a+b)} z_1$, and by Lemma 6.2, as z moves from z_1

to z_2 on half of the circle C_{ab} that contains x , $f(z) =_{\text{def}} \|z - c\|$ ($z \in C_{ab}$) increases, and since it reaches the value 1 at $z = x$ ($\|x - c\| = 1$, since $c \in C_{xy}$), $f(z) = \|z - c\| > \|x - c\| = 1$ when z lies on the short arc $\text{arc}(x, z_2)$ of C_{ab} . A similar inequality holds when z moves from z_1 to z_2 on half of the circle C_{ab} that contains y . This shows that $\|z - c\| > 1$ for $z \in C_{ab} \setminus A_{xy}$ (and $c \in C_{xy} \setminus A_{ab}$). ■

Proof of Theorem 8.1: The edge e (whose vertices are a, b) is a short arc of the circle C_{xy} , by Corollary 7.2 above. Hence $e = A_{ab}$, where A_{ab} and A_{xy} are the short circular (dual) arcs defined for the couple of 2-tuples $(a, b; x, y)$ in Definition 8.2 above. Put $H =_{\text{def}} \text{aff}(x, a, b)$ and let H^+ be the closed half space bounded by H which contains y . Since A_{xy} is a short arc of C_{ab} , clearly $A_{xy} \subset H^+$.

Claim 1: $(\text{relint} A_{xy}) \cap V = \emptyset$.

Proof: If A_{xy} contains a point $v \in V$ in its relative interior, then by Lemma 8.1 a) above $B(v, 1) \cap \text{relint} A_{ab} = \emptyset$, hence $e = A_{ab} \not\subset \mathcal{B}(V)$, which is a contradiction (e is an edge of $\mathcal{B}(V)$). This proves Claim 1.

Claim 2: *There is an edge of $\mathcal{B}(V)$ incident with x whose initial arc from x lies on A_{xy} .*

Proof: Put $m =_{\text{def}} \#(\text{vert} F_x)$. If $m = 2$, i.e., $\text{vert} F_x = \{a, b\}$, then by Lemma 6.4 d) there are two edges of $\mathcal{B}(V)$ incident with x which have initial arcs from x on C_{ab} , lying in different sides of H ; hence one of them has an initial arc from x lying on $A_{xy} (\subset C_{ab})$ contained in H^+ . If $m \geq 3$ then, by Proposition 6.4 (iv), the point y and $(\text{vert} F_x) \setminus \{a, b\}$ lie on the same side of the plane H , and since $y \in H^+$, $(\text{vert} F_x) \setminus \{a, b\} \subset H^+$. By Lemma 6.4 c) there is an edge e' of $\mathcal{B}(V)$ that has in initial arc from x lying on C_{ab} contained in H^+ , i.e., the initial arc of e' from x is contained in A_{xy} . This proves Claim 2.

Claim 3: $A_{xy} = e'$.

Proof: e' has an initial arc from x lying on A_{xy} (Claim 2), hence its other endpoint (in $\text{int} H^+$) is either y or a point $y' \in \text{relint} A_{xy}$. The second case is impossible by Claim 1. This proves Claim 3, concluding the proof of Theorem 8.1. ■

The following lemma is needed in the proof of Theorem 8.2 below.

Lemma 8.2: *Let $V \subset \mathbb{R}^3$ be an extremal set for the Vaszónyi problem with $\infty > \#V = n \geq 4$, and e, f be two dual edges of $\mathcal{B}(V)$ with endpoints (x, y) and (a, b) , respectively, and let $u \in \text{relint} e$. Then:*

- a) $V \cup \{u\}$ is an extremal set.
- b) All edges of $\mathcal{B}(V)$ are edges of $\mathcal{B}(V \cup \{u\})$ and vice versa, except for e and f , for which
 - (i) the edge e of $\mathcal{B}(V)$ is splitted in $\mathcal{B}(V \cup \{u\})$ into two edges e_1, e_2 with endpoints (x, u) and (u, y) , respectively, and
 - (ii) $\mathcal{B}(V \cup \{u\}) \cap \text{relint} f = \emptyset$, and instead of the edge f of $\mathcal{B}(V)$, $\mathcal{B}(V \cup \{u\})$ has two new edges f_1, f_2 which are the duals of the two edges e_1, e_2 to which e was splitted, respectively.

Proof:

- a) e is an arc of the circle $C_{ab} = S_a \cap S_b$, hence $\|a - u\| = \|b - u\| = 1$, and since e is an edge of $\mathcal{B}(V)$ and $u \in \text{relint} e$, u is not a vertex of $\mathcal{B}(V)$, hence $\|u - z\| < 1$ for $z \in V \setminus \{a, b\}$. Thus $\text{diam}(V \cup \{u\}) = 1$ and $e(V \cup \{u\}) = e(V) + 2 = (2n - 2) + 2 = 2(n + 1) - 2 = e(n + 1, 3)$ (see Corollary 7.1 above), i.e., $V \cup \{u\}$ is extremal.
- b) By Lemma 8.1 a) we have $\|u - z\| > 1$ for $z \in \text{relint} f$, i.e., $B(u) \cap \text{relint} f = \emptyset$, hence $\mathcal{B}(V \cup \{u\}) \cap \text{relint} f = \emptyset$ and f is not an edge of $\mathcal{B}(V \cup \{u\})$. Now u is a dangling vertex of $\mathcal{B}(V \cup \{u\})$, hence the edge e of $\mathcal{B}(V)$ is splitted by u into two (new) edges e_1, e_2 of $\mathcal{B}(V \cup \{u\})$. Clearly, their dual edges in $\mathcal{B}(V) \cup \{u\}$, namely f_1 and f_2 , respectively, in $\mathcal{B}(V \cup \{u\})$ are not edges of $\mathcal{B}(V)$, i.e., they are also new edges relatively to $\mathcal{B}(V)$. It remains to show that every edge g of $\mathcal{B}(V)$ other than e and f is also an edge of $\mathcal{B}(V \cup \{u\})$. It suffices to show that $\text{relint} g \subset \text{int} B(u)$, i.e., that the following claim holds.

Claim: $\|u - d\| < 1$ for $d \in \text{relint} g$.

Proof: Let (s, t) be the two endpoints of g and consider two cases:

Case I: $\{s, t\} = \{a, b\}$, i.e., g and f share the same endpoints. Let h be the dual edge of g in $\mathcal{B}(V)$. Clearly, h is a short arc of the circle C_{ab} , same as e (which is the dual arc of f), so both e and h lie on C_{ab} , and clearly $\text{relint } e \cap \text{relint } h = \emptyset$. Let x', y' be the endpoints of h ($x', y' \in C_{ab}$).

It follows that u , as a point from the relative interior of $e \subset C_{ab}$, lies on the *long* arc between x' and y' of the circle C_{ab} . Since (a, h, x', y') are in equilateral position (Definition 8.2), we can apply Lemma 8.1 a) to the dual arcs $g = A_{ab} \subset C_{x'y'}$ and $h = A_{x'y'} \subset C_{ab}$, with $c = u \in C_{ab} \setminus A_{x'y'} = C_{ab} \setminus h$ and $z = d \in \text{relint } A_{ab} = \text{relint } g$, to conclude that $\|z - d\| < 1$ ($\|c - z\| < 1$ in the notations of Lemma 8.1 a)).

Case II: $\{s, t\} \neq \{a, b\}$. Then $\|u - s\| < 1$ or $\|u - t\| < 1$ (or both), say $\|u - s\| < 1$. Assume, by r.a.a., that there exists a $d \in \text{relint } g$ such that $\|u - d\| > 1$. Then there is a point $z \in \text{relint } g$ on the part of g between s and d such that $\|u - z\| = 1$. Then $\|z - a\| = \|z - b\| = \|z - u\| = 1$, and $z \in \text{bd } \mathcal{B}(V \cup \{u\})$. Thus $z \in \text{vert } \mathcal{B}(V \cup \{u\})$. On the other hand, $z \in \text{relint } g$, hence $z \notin \text{vert } \mathcal{B}(V) = V$ (the equality holds by Theorem 7.1 (iii) – the extended GHS-Theorem). It follows that $z \notin V \cup \{u\}$, hence $V \cup \{u\} \neq \text{vert } \mathcal{B}(V \cup \{u\})$. Since $V \cup \{u\}$ is extremal (part a)) this is a contradiction to Theorem 7.1 (iii), proving the Claim. ■

The following theorem shows how to obtain an extremal set $V \cup W$ from a given extremal set V by adding a set W of “dangling vertices” at will – just observing the simple rule that no two new points belong to dual edges of $\mathcal{B}(V)$.

Theorem 8.2: *Let $V \subset \mathbb{R}^3$ be an extremal set for the Vaszónyi problem with $\infty > \#V = n \geq 4$. Let $W \subset \text{skel}_1 \mathcal{B}(V)$, where W is finite such that for any two dual edges e, f of $\mathcal{B}(V)$ either $W \cap e = \emptyset$ or $W \cap f = \emptyset$. Then $V \cup W$ is extremal.*

Proof: Let $W = \{v_1, v_2, \dots, v_m\}$, $1 \leq m < \infty$, and let e_i be the edge of $\mathcal{B}(V)$ that contains v_i , $1 \leq i \leq m$. Assume, w.l.o.g., that $v_i \in \text{relint } e_i$ ($1 \leq i \leq m$). By assumption there are no two dual edges among $e_1, e_2, \dots, e_m$. Put $V_0 =_{\text{def}} V$, $V_1 =_{\text{def}} V_0 \cup \{v_1\}$, $V_2 =_{\text{def}} V_1 \cup \{v_2\}$, $\dots$, $V_m = V_{m-1} \cup \{v_m\} = V \cup W$.

Claim: V_i is an extremal set for $0 \leq i \leq m$, and e_{i+1} is an edge of $\mathcal{B}(V_i)$ for $0 \leq i \leq m-1$.

Remark: The part of the claim that e_{i+1} is an edge of $\mathcal{B}(V_i)$ is inserted only for technical reasons: to carry on the induction step (see below).

For $i = 0$ there is nothing to prove.

Induction step $i \longrightarrow i + 1$, $0 \leq i < m$:

Assume that V_i is extremal and that e_{i+1} is an edge of $\mathcal{B}(V_i)$. Consider $V_{i+1} = V_i \cup \{v_{i+1}\}$. Since $v_{i+1} \in \text{relint } e_{i+1}$ and e_{i+1} is an edge of $\mathcal{B}(V_i)$, $V_{i+1} = V_i \cup \{v_{i+1}\}$ is extremal, by Lemma 8.2 a). By assumption the edge e_{i+2} is not dual to e_{i+1} (if $i \leq m - 2$; if $i = m - 1$, this part of the induction proof is finished), and we conclude from Lemma 8.2 b) that e_{i+2} is an edge of $\mathcal{B}(V_{i+1})$ as well. This proves the claim.

It follows from the claim that, in particular, $V_m = V \cup W$ is extremal. This proves Theorem 8.2. ■

§ 9. Fixed-point free property of φ (strong self-duality)

In this paragraph we define the first barycentric subdivision of $\mathcal{SF}(\mathcal{B}(V))$ for an extremal set V and show that the canonical self-duality φ of $\mathcal{SF}(\mathcal{B}(V))$ is fixed-point free when it acts as an automorphism of this barycentric subdivision.

Let $V \subset \mathbb{R}^3$ be an extremal set with $\#V = n \geq 4$. We are about to define the *barycentric subdivision* of the spherical complex $\mathcal{SF}(\mathcal{B}(V))$. This will be done in two stages. First we define an abstract two-dimensional simplicial complex $\mathcal{K}$. Then we define a (curvilinear) realization of $\mathcal{K}$ as a subdivision of $\mathcal{SF}(\mathcal{B}(V))$.

I. Vertices of $\mathcal{K}$: Corresponding to each non-empty face Φ of $\mathcal{SF}(\mathcal{B}(V))$, $\mathcal{K}$ has a vertex $z(\Phi)$. Thus we have vertices $z(x)$ for vertices $x \in V (= \text{vert } \mathcal{B}(V))$, by Theorem 7.1 (iii), vertices $z(e)$ for edges e of $\mathcal{SF}(\mathcal{B}(V))$, and vertices $z(F)$ for facets $F = F_x$ of $\mathcal{SF}(\mathcal{B}(V))$. The correspondence $\Phi \rightarrow z(\Phi)$ is 1-1.

II. Faces of $\mathcal{K}$: For each *chain* (*flag* in the current terminology) $\mathcal{C}$ of faces of $\mathcal{SF}(\mathcal{B}(V))$ the set $\{z(\Phi) : \Phi \in \mathcal{C}\}$ is a *face* of $\mathcal{K}$.

The empty chain yields the empty face $\emptyset$ of $\mathcal{K}$. Chains of size 1 yield faces consisting of single vertices of $\mathcal{K}$. The edges of $\mathcal{K}$ (1-faces) are of three kinds:

- $\{z(x), z(e)\}$, where e is an edge of $\mathcal{SF}(\mathcal{B}(V))$, and $x \in V$ is an endpoint of e ;
- $\{z(e), z(F)\}$, where F is a facet of $\mathcal{SF}(\mathcal{B}(V))$, and e is an edge of F ;
- $\{z(x), z(F)\}$, where F is a facet, and x is a vertex of F .

The 2-faces of $\mathcal{K}$ are of the form $\{z(x), z(e), z(F)\}$, where F is a facet of $\mathcal{SF}(\mathcal{B}(V))$, e is an edge of F , and $x \in V$ is an endpoint of e .

Now we present the realization. We start with the *vertices* of $\mathcal{K}$. For each vertex $x \in V$ of $\mathcal{SF}(\mathcal{B}(V))$, we realize $z(x)$ by x .

For each edge e of $\mathcal{SF}(\mathcal{B}(V))$, we realize $z(e)$ by the midpoint of the circular arc e . Each facet $F = F_y$ of $\mathcal{SF}(\mathcal{B}(V))$, $y \in V$, is a compact, strictly spherically convex subset of the unit sphere $S(y, 1)$ (cf. Corollary 5.2 above). We realize $z(F)$ by a point that lies in the relative interior of the spherical convex hull of the vertices of F on $S(y, 1)$. (When F is a digon, then the spherical convex hull of its vertices is just the geodesic arc on $S(y, 1)$ that connects the two vertices.)

Now we come to the *edges* of $\mathcal{K}$. The edge $\{z(x), z(e)\}$ (where e is an edge of $\mathcal{SF}(\mathcal{B}(V))$, and x is an endpoint of e) is realized by the half of e with endpoints $x, z(e)$. The edge $\{z(x), z(F_y)\}$ is realized by the geodesic arc on $S(y, 1)$ with endpoints $z(x)$ and $z(F_y)$. Similarly, the edge $\{z(e), z(F_y)\}$ is realized by the geodesic arc on $S(y, 1)$ with endpoints $z(e)$ and $z(F_y)$.

We continue with the *facets* of $\mathcal{K}$. The facet $\{z(x), z(e), z(F_y)\}$ is realized by the union of all geodesic arcs on $S(y, 1)$ that connect $z(F_y)$ with points on the half of e with endpoints x and $z(e)$. This is the closed convex triangular region on $S(y, 1)$ bounded by the realization of the edges of $\{z(x), z(e), z(F_y)\}$.

To make the correspondence between (a rectilinear realization of) $\mathcal{K}$ and the curvilinear realization described above more explicitly, we introduce the following notation: If C is a circular arc with endpoints a and b , on a circle with center c , we denote by “ $1 - \lambda$ ” a +“ λ ” b ($0 \leq \lambda \leq 1$) the point x on C

that satisfies $\nabla acx = \lambda \nabla acb$. If $p = \alpha z(F_y) + \beta z(e) + \gamma z(x)$ is a point in the triangle $[z(F_y), z(e), z(x)]$ of $\mathcal{K}$ ($x \in e, e \in F_x, \alpha, \beta, \gamma \geq 0, \alpha + \beta + \gamma = 1$), then the corresponding point p' in the curvilinear realization is defined in two steps as follows:

$$q' = \frac{\beta}{1-\alpha} z(e) + \frac{\gamma}{1-\alpha} z(x),$$

where q' lies on e , between $z(x)$ and $z(e)$,

$$p' = \alpha z(F_y) + (1-\alpha) q',$$

where p' lies on the great circle through $z(F_y)$ and q' on $S(y, 1)$.

The canonical self-duality φ of $\mathcal{SF}(\mathcal{B}(V))$ induces an automorphism (to be denoted again by φ) of order 2 on the barycentric subdivision $\mathcal{K}$. To show that this automorphism is fixed-point free, it is sufficient (and also necessary) to confirm that no non-empty cell of $\mathcal{K}$ is mapped to itself.

Indeed, if a cell C of $\mathcal{K}$ contains $z(x), x \in V$, then $\varphi(C)$ contains $\varphi(z(x)) = z(F_x)$ and, hence, does not contain $z(x)$, since x is *not* a vertex of F_x , and therefore $\varphi(C) \neq C$. Similarly, if $z(F_x) \in C$, then $z(x) \in \varphi(C)$, and therefore $z(F_x) \notin \varphi(C)$, hence $\varphi(C) \neq C$. There remains the case where C consists of a single vertex $z(e)$ for some edge of $\mathcal{SF}(\mathcal{B}(V))$. But then $\varphi(C) = \{z(e')\}$, where e' is the edge dual to e . Clearly, $e' \neq e$, hence $\varphi(C) \neq C$ also in this case.

One can easily check that an even stronger statement holds: If C is a cell of $\mathcal{K}$, then C and $\varphi(C)$ have no vertex in common. We sum the foregoing discussion as


Theorem – Definition 9.1 (strong self duality): *The canonical (involutory) self-duality $\varphi : \mathcal{SF}(\mathcal{B}(V)) \rightarrow \mathcal{SF}(\mathcal{B}(V))$ for V extremal is fixed point free. An involutory self-duality of a 2-dimensional polyhedral (maybe curvilinearly realized as $\mathcal{SF}(\mathcal{B}(V))$ above) complex which is fixed point free is a strong self-duality. Thus $\mathcal{SF}(\mathcal{B}(V))$ is strongly self-dual.*

The following example, illustrating the notion of self-dual complex introduced above where the complex is that of the boundary of a 3-polytope, is a generalization of an example given by Grünbaum and Shephard [15].

Example 9.1 (An apexed 3-prism as a 3-polytope with many involutory self-dualities and only one strong self duality):

Let P be a 3-polytope which is the union of a 3-prism with (parallel) bases Q, Q' and a 3-pyramid with basis Q' apexed at q , such that $\text{aff}Q'$ separates Q and q (equivalently: $\text{relint}[q, v] \cap \text{relint}Q' \neq \emptyset$ for $v \in \text{vert}Q$). Note that Q is a facet of P and Q' is not a facet. The polytope P is an *apexed prism* (of dimension 3).

Put $n =_{\text{def}} \#\text{vert}Q (= \#\text{vert}Q'), n \geq 3$, and denote by $a_1, \dots, a_n$, resp. $b_1, \dots, b_n$ the vertices of Q , resp. Q' , ordered cyclically on $\text{relbd}Q$, resp. $\text{relbd}Q'$. Assume, w.l.o.g., that $[a_i, b_i]$ is an edge of P for $1 \leq i \leq n$ (otherwise shift (cyclically) the indices of the b_i 's and, if necessary, invert their order).



– Figure 13 –

The 2-dimensional face structure $\mathcal{F}(P)$ of P can be drawn as a Schlegel diagram on the facet Q (see Figure 13 for $n = 6$; ignore the A_i 's and B_i 's meanwhile). Assume, w.l.o.g., that q lies “above” the plane of the figure and that, looking on the diagram from the side of q (i.e., from “above”), the ordering of $a_1, \dots, a_n$, resp. $b_1, \dots, b_n$, is clockwise. Note that a_i is 3-valent and b_i is 4-valent in $\text{skel}_1(P)$.

We will describe now all the involutory self-dualities of $\mathcal{F}(P)$. Assume that $\psi : \mathcal{F}(P) \rightarrow \mathcal{F}(P)$ is an involutory self-duality of $\mathcal{F}(P)$. Put $A_i = \psi(a_i)$ and $B_i = \psi(b_i), 1 \leq i \leq n$.

Claim 1: q is a vertex of A_i for $1 \leq i \leq n$.

Proof: a_i is 3-valent in $\text{skel}_1 P$, hence $\psi(a_i)$ is a triangular facet. If $n \geq 4$, then the facets around q are the only triangular facets of P , hence $\psi(a_i) = A_i$ must have q as a vertex. For $n = 3$ an extra consideration is needed, because then Q being a triangular facet is also a candidate for $A_i = \psi(a_i)$ for some i . Assume, by r.a.a., that $\psi(a_i) = Q$ for some $1 \leq i \leq 3$, say, w.l.o.g., $\psi(a_1) = Q$. Then $\psi(a_2)$ (and $\psi(a_3)$) must have q as a vertex and the edge $[a_1, a_2]$ of P is mapped by ψ to the edge $\psi(a_1) \cap \psi(a_2) = Q \cap \psi(a_2)$. This is a contradiction, since Q and a facet having q as a vertex do not share an edge (in fact, they have an empty intersection).

Claim 2: $q \xleftrightarrow{\psi} Q$.

Proof: Since a_i , $1 \leq i \leq n$, are the vertices of Q , their images $\psi(a_i) = A_i$ must all share $\psi(Q)$ as a vertex. But A_i , $1 \leq i \leq n$, share only q by Claim 1, hence $\psi(Q) = q$.

Claim 3: B_i ($1 \leq i \leq n$) and Q share an edge.

This follows from Claims 1 and 2 above.

Claim 4: q is a vertex of the edge $[a_i, a_{i+1}] \xleftrightarrow{\psi} A_i \cap A_{i+1}$ ($1 \leq i \leq n$).

Proof: $[a_i, a_{i+1}]$ is an edge, hence its image by ψ , $A_i \cap A_{i+1}$, is an edge, and by Claim 1 all A_i 's share q as a vertex. Thus also Claim 4 is verified.

It follows from Claim 4 that the facets $A_1, \dots, A_n$ are cyclically ordered around q , either clockwise (same as the a_i 's) or counterclockwise. Since $[a_1, b_1]$ is an edge, its image under ψ is the edge $\psi(a_1) \cap \psi(b_1) = A_1 \cap B_1$. Since $A_i \cap B_i$ has to be of the form $[a_j, a_{j+1}]$ for some $1 \leq j \leq n$ the edge $A_1 \cap B_1$ is of the form $[b_\nu, b_{\nu+1}]$ for some $1 \leq \nu \leq n$ ($n = 6$ and $\nu = 4$ in Figure 16 above). It follows that either

(I) $A_i \cap B_i = [b_{\nu-i+1}, b_{\nu-i+2}]$, $1 \leq i \leq n$, or

(II) $A_i \cap B_i = [b_{\nu+i-1}, b_{\nu+i}]$, $1 \leq i \leq n$.

We discuss these two cases separately.

Case (I): In this case the A_i 's and B_i 's ($1 \leq i \leq n$) are cyclically ordered counterclockwise around q (*opposite* to the cyclic order of the a_i 's and b_i 's).

Clearly, in this case

$$\begin{aligned} (1) \quad & B_i \cap B_{i+1} = [a_{\nu-i+1}, b_{\nu-i+1}], \\ (2) \quad & A_i \cap A_{i+1} = [q, b_{\nu-i+1}], \\ (3) \quad & Q \cap B_i = [a_{\nu-i+1}, a_{\nu-i+2}]. \end{aligned} \tag{36}$$

Since $i = \nu - i + 1 \Leftrightarrow \nu$ is odd and $i = \frac{\nu+1}{2}$, and $i = \nu - i + 2 \Leftrightarrow \nu$ is even and $i = \frac{\nu+2}{2}$, it follows from (I) that

$$b_i \in B_i \Leftrightarrow \begin{cases} i = \frac{\nu+1}{2} & \text{if } \nu \text{ is odd,} \\ i = \frac{\nu+2}{2} & \text{if } \nu \text{ is even.} \end{cases} \tag{37}$$

Thus in case (I), (36) + (I) give four necessary conditions for ψ to be an involutory self-duality of $\mathcal{F}(P)$.

Claim 5: *The conditions (36) + (I) are (necessary and) sufficient for ψ to be an involutory self duality of $\mathcal{F}(P)$.*

Proof: With the notation of q, Q, a_i, b_i, A_i, B_i ($1 \leq i \leq n$) and ν ($1 \leq \nu \leq n$) satisfying (I) + (36) introduced above, define an involutory self-duality $\psi_\nu : \mathcal{F}(P) \rightarrow \mathcal{F}(P)$ in two stages. First define ψ_ν on the vertices and facets of $\mathcal{F}(P)$ by $q \xleftrightarrow{\psi_\nu} Q$, $a_i \xleftrightarrow{\psi_\nu} A_i$, $b_i \xleftrightarrow{\psi_\nu} B_i$ ($1 \leq i \leq n$).

In order to define ψ_ν on the edges of $\mathcal{F}(P)$, observe first that there are four types of edges and that each edge has two forms, as a segment with two endpoints and dually as an intersection of two neighboring facets (in the dual graph of $\mathcal{F}(P)$). Here is the list of these four types, each edge is given by its two dual forms (based on (I) + (36)).

$$\begin{aligned} 1) \quad & [a_i, a_{i+1}] = Q \cap B_{\nu-i+1}, \\ 2) \quad & [a_i, b_i] = B_{\nu-i+1} \cap B_{\nu-i+2}, \\ 3) \quad & [b_i, b_{i+1}] = A_{\nu-i+1} \cap B_{\nu-i+1}, \\ 4) \quad & [q, b_i] = A_{\nu-i+1} \cap A_{\nu-i+2}. \end{aligned} \tag{38}$$

For each edge we use its left hand side form in (38) to define its image by ψ_ν .

$$\begin{aligned}
1) \quad [a_i, a_{i+1}] &\xleftrightarrow{\psi_\nu} A_i \cap A_{i+1}, \\
2) \quad [a_i, b_i] &\xleftrightarrow{\psi_\nu} A_i \cap B_i, \\
3) \quad [b_i, b_{i+1}] &\xleftrightarrow{\psi_\nu} B_i \cap B_{i+1}, \\
4) \quad [q, b_i] &\xleftrightarrow{\psi_\nu} Q \cap B_i.
\end{aligned} \tag{39}$$

In order to show that ψ_ν is well defined on the edges, we must check that the following four diagrams are commutative.

$$\begin{aligned}
1) \quad [a_i, a_{i+1}] &= Q \cap B_{\nu-i+1} \\
&\quad \begin{array}{ccc} & \nwarrow \psi_\nu & \nearrow \psi_\nu \\ & A_i \cap A_{i+1} & \end{array} \\
2) \quad [a_i, b_i] &= B_{\nu-i+1} \cap B_{\nu-i+2} \\
&\quad \begin{array}{ccc} & \nwarrow \psi_\nu & \nearrow \psi_\nu \\ & A_i \cap B_i & \end{array} \\
3) \quad [b_i, b_{i+1}] &= A_{\nu-i+1} \cap B_{\nu-i+1} \\
&\quad \begin{array}{ccc} & \nwarrow \psi_\nu & \nearrow \psi_\nu \\ & B_i \cap B_{i+1} & \end{array} \\
4) \quad [q, b_i] &= A_{\nu-i+1} \cap A_{\nu-i+2} \\
&\quad \begin{array}{ccc} & \nwarrow \psi_\nu & \nearrow \psi_\nu \\ & Q \cap B_i & \end{array}
\end{aligned}$$


Let us check this for diagram 1). The left double arrow in 1) follows from (39), 1). To show the right double arrow, write

$$\begin{aligned}
Q \cap B_{\nu-i+1} &\xleftrightarrow{\psi_\nu} [q, b_{\nu-i+1}] \text{ (by (39), 4))} \\
&= A_{\nu-(\nu-i+1)+1} \cap A_{\nu-(\nu-i+1)+2} \text{ (by (38), 4))} \\
&= A_i \cap A_{i+1}, \text{ i.e.,} \\
Q \cap B_{\nu-i+1} &\xleftrightarrow{\psi_\nu} A_i \cap A_{i+1}.
\end{aligned}$$

Similarly one can check the commutativity of the remaining diagrams 2), 3), and 4). This shows that ψ_ν ($1 \leq \nu \leq n$) is an involutory self-duality, proving Claim 5. $\blacksquare$

By (37), ψ_ν has a fixed point (when acting as an automorphism of the first barycentric subdivision of $\mathcal{F}(P)$): the edge corresponding to the flag (b_i, B_i) , where $b_i \in B_i$ ($i = \frac{\nu+1}{2}$, resp. $i = \frac{\nu+2}{2}$, for ν odd, resp. even), is mapped by ψ_ν to itself (since $b_i \xleftrightarrow{\psi_\nu} B_i$). Thus all the involutory self-dualities ψ_ν , $1 \leq \nu \leq n$, are *not* strong self-dualities.

Case (II): In this case the A_i 's and B_i 's are cyclically ordered clockwise around q (*same* as the cyclic order of the a_i 's and b_i 's).



– Figure 14 –

Figure 14 sketches the Schlegel diagram of P in this case. We have now

$$[a_1, b_1] = B_{n-\nu+1} \cap B_{n-\nu+2} \text{ and } [a_1, b_1] \xleftrightarrow{\psi} A_1 \cap B_1 = [b_\nu, b_{\nu+1}],$$

hence,

$$[b_\nu, b_{\nu+1}] \xleftrightarrow{\psi} B_{n-\nu+1} \cap B_{n-\nu+2} \xleftrightarrow{\psi} [b_{n-\nu+1}, b_{n-\nu+2}].$$

It follows that $[b_\nu, b_{\nu+1}] = [b_{n-\nu+1}, b_{n-\nu+2}]$, hence $b_\nu = b_{n-\nu+1}$, i.e., $\nu = n - \nu + 1$, i.e.,

$$\nu = \frac{n+1}{2}. \quad (40)$$

Thus case (II) is possible only if n is odd and $\nu = \frac{n+1}{2}$ (unlike case (I) in which all ν 's, $1 \leq \nu \leq n$, are possible). In this case we have, in addition to (II), the following three conditions (compare to conditions (I) + (36) above):


$$\begin{aligned} 1) \quad B_i \cap B_{i+1} &= [a_{\nu+i}, b_{\nu+i}], \\ 2) \quad A_i \cap A_{i+1} &= [q, b_{\nu+i}], \\ 3) \quad Q \cap B_i &= [a_{\nu+i-1}, a_{\nu+i}]. \end{aligned} \quad (41)$$

Thus the conditions

$$n \geq 3 \text{ is odd, } \nu = \frac{n+1}{2}, \text{ (II), and (41)} \quad (42)$$

are necessary for ψ to be an involutory self-duality of $\mathcal{F}(P)$.

Claim – Definition 6: *The conditions (42) are (necessary and) sufficient for ψ to be an involutory self-duality of $\mathcal{F}(P)$. Moreover, these conditions uniquely determine ψ . We denote ψ , uniquely determined thus by ψ^* .*



– Figure 15 –

Proof: The proof is similar to that of Claim 5 above. One has to write four equalities like (38), define $\psi^* : \mathcal{F}(P) \rightarrow \mathcal{F}(P)$ similarly to the definition of ψ_ν in (39), and then to check the commutativity of four diagrams as above. We kindly ask the reader to make this check. Figure 15 illustrates the case $n = 5$ ($\nu = \frac{5+1}{2} = 3$).

$\psi^* : \mathcal{F}(P) \rightarrow \mathcal{F}(P)$ is defined by $q \xleftrightarrow{\psi^*} Q$, $a_i \xleftrightarrow{\psi^*} A_i$, and $b_i \xleftrightarrow{\psi^*} B_i$ ($1 \leq i \leq 5$), and on the edges ψ^* is defined as ψ_ν in (39).

Claim 7: ψ^* (assured by Claim and Definition 6 above) is fixed point free (as an induced automorphism of the first barycentric subdivision of $\mathcal{F}(P)$).

Proof: Again we leave this for the reader, using the equalities and definition of ψ^* as in his proof of Claim and Definition 6 above. Alternatively, the reader may check this directly for the case $n = 5$ ($\nu = 3$) depicted in Figure 15 above. ■

Here is a summary of the foregoing discussion:

Proposition 9.1: *Let P be an apexed 3-prism with (polygonal) basis of order $n \geq 3$.*

- a) *The face complex $\mathcal{F}(P)$ of P has exactly n involutory self-dualities ψ_ν , $\nu = 1, 2, \dots, n$, each having a fixed point (in the sense of the barycentric subdivision given above).*
- b) *For $n \geq 3$ odd, $\mathcal{F}(P)$ has a unique fixed point free involutory self-duality, and for $n > 3$ even it has none.*
- c) *Hence $\mathcal{F}(P)$ is strongly self-dual iff $n \geq 3$ is odd.*

From Proposition 9.1 c) we get

Corollary 9.1: *There is no extremal set $V \subset \mathbb{R}^3$, $4 \leq \#V < \infty$, such that $\mathcal{SF}(\mathcal{B}(V))$ is isomorphic to the face complex $\mathcal{F}(P)$ of an apexed prism P whose basis is of an even order $n \geq 4$.*

In view of Corollary 9.1, knowing that $\mathcal{SF}(\mathcal{B}(V))$ is strongly self dual for an extremal set V , a natural question arises. Is there an extremal set $V \subset \mathbb{R}^3$ such that $\mathcal{SF}(\mathcal{B}(V))$ is combinatorially equivalent to the face complex of an

apexed prism whose basis is of an *odd* order $n \geq 3$? The answer is “yes”, as the following example shows.

Example 9.2 (an extremal configuration the face complex of whose ball polytope is combinatorially equivalent to that of an apexed 3-prism): Let $V = \{c\} \cup \text{vert} P_{2k-1}$ ($k \geq 2$) be the set of vertices of the suspended $(2k-1)$ -gon described in Example 1.2 above. $\mathcal{SF}(\mathcal{B}(V))$ is combinatorially equivalent to the face complex $\mathcal{F}(\Pi)$ of a 3-pyramid Π whose basis is of order $2k-1$, as one can easily check. Put $p_c =_{\text{def}} c - \frac{c}{\|c\|}$; p_c is the central point of the facet F_c of $\mathcal{B}(V)$ ($\|p_c - c\| = 1$), and let p'_c be a point such that $p'_c =_{\text{def}} p_c + \varepsilon \|p_c - c\|$ for some $\varepsilon > 0$. Note that $p'_c \notin \mathcal{B}(V)$ and that p'_c lies “beyond” the facet F_c , relatively to c . If ε is small enough ($\varepsilon \leq 0.1$ will certainly suffice), then $S(p'_c) \cap \text{skel}_1 \mathcal{B}(V)$ (where $S(p'_c)$ is the unit sphere centered at p'_c) consists of $2k-1$ points lying on the $2k-1$ edges of $\mathcal{B}(V)$ incident with c , respectively, all of which are very close to c (certainly of distance not more than 3ε apart from c for $\varepsilon \leq 0.1$). Denote this set of $2k-1$ “new” points by W_{2k-1} and define $W =_{\text{def}} \{p'_c\} \cup \text{vert} P_{2k-1} \cup W_{2k-1}$ or, equivalently, $W =_{\text{def}} \{p'_c\} \cup (\text{vert} \mathcal{B}(V) \setminus \{c\}) \cup W_{2k-1}$. Clearly, $\#W = 2n+1$, where $n =_{\text{def}} 2k-1$ (odd) and $\text{diam } W = 1$.

Claim: W is an extremal configuration for the Vázsonyi problem.

Proof: There are n edges of $D(W)$ in $D(\text{vert} P_{2k-1})$, n edges in $D(p'_c \cup W_{2k-1})$, and $2n$ edges with one endpoint in $\text{vert} P_{2k-1}$ and one endpoint in W_{2k-1} (since every vertex of P_{2k-1} is incident with two points in W_{2k-1}). Hence $e(W) = n + n + 2n = 4n = 2(2n+1) - 2 = e(2n+1, 3)$, i.e., W is extremal. This proves the claim.

It follows that $\mathcal{SF}(\mathcal{B}(W))$ is strongly self-dual (via the canonical self-duality φ). Moreover, $\mathcal{SF}(\mathcal{B}(V))$ is isomorphic to the face complex $\mathcal{F}(P)$ of an apexed 3-prism P whose basis is of order $n (= 2k-1)$. We leave this check to the reader with just one hint: p'_c plays the role of the apex of P , and $F_{p'_c}$ plays the role of its basis. Clearly, the canonical self-duality φ of $\mathcal{SF}(\mathcal{B}(V))$ corresponds to the strong self-duality of $\mathcal{F}(P)$ whose existence and uniqueness is assured by Proposition 9.1 b) above.

Remarks 9.1:

- 1) As far as we know, the configuration W just now described above is a

completely new type of extremal configuration for the Vázsonyi problem. Moreover, although the foregoing construction of W was rigidly described, it allows a considerable degree of freedom. First, the initial set V of the suspended $(2k - 1)$ -gon allows some perturbations of $\text{vert}P_{2k-1}$ on S_c , as hinted in Example 1.2 above. Second, the point p'_c , i.e., the center of the unit ball $\mathcal{B}(p'_c)$ by which c was “truncated” from $\mathcal{B}(V)$ to obtain $\mathcal{B}(W)$, allows some perturbations near p_c as well.

- 2) In fact, this idea of ball truncation is the germ of a vastly general new construction of extremal configurations W given an extremal configuration V : choose a vertex $c \in \text{vert}\mathcal{B}(V)$ ($= V$ by Theorem 7.1 (iii)), choose a point $p_c \in \text{relint}F_c$, move p_c slightly out of $\mathcal{B}(V)$ to obtain a point p'_c lying beyond the facet F_c relatively to c , and put $W =_{\text{def}} \{p'_c\} \cup (\mathcal{B}(V) \cap S(p'_c)) \cup (V \setminus \{c\})$. The following is easy to prove.

Claim: *If p'_c is close enough to p_c , then $\text{diam}W = 1$ and W is extremal.*

In a subsequent paper we will exploit the idea of ball truncation and its variants, e.g., that p'_c may lie not only beyond a facet but also beyond an edge of $\mathcal{B}(V)$, and even beyond a vertex. Meanwhile we refer to our paper [21], where this construction is explained in a more detailed way.

Remarks 9.2: Applications to strongly self-dual 3-polytopes.

- 1) The “ball truncation” described above has direct and obvious analogues in the construction of a strongly self-dual 3-polytope Q (abbr. *SSD* 3-polytope) given a *SSD* 3-polytope Q . (*SSD* means “Strongly Self-Dual” in the sense of **Theorem – Definition 9.1** above.) This rich family of 3-polytopes seems to be an uncharted territory, and we hope that the present work will prompt a study of it. One aim of such a study might be to construct many (if not all) of the *SSD* 3-polytopes by using analogues of the ball truncation constructions described above. The model we have in mind for such a study is Steinitz’s proof that every planar 3-connected graph is polytopal (“Steinitz’s theorem”) as it appears in [14], pp. 235–243. This proof shows in fact that every 3-polytope P can be obtained from the tetrahedron by a series of either truncations of 3-valent vertices or edges or triangular facets (the

opposite operations of the “ η_i -reductions” in Fig. 13.1.2 in p. 238 there) or addition of a point beyond a triangular facet (the opposite ω_i -reductions in Fig. 13.1.1 in p. 237 there).

- 2) Sallee asks in [25], p. 322, two questions which in the language of the present work can be formulated as follows.

Question 1: Is the face structure $\mathcal{SF}(\mathcal{B}(V))$ of every Ball polytope $\mathcal{B}(V)$ of an extremal set $V \subset \mathbb{R}^3$ ($4 \leq \#V < \infty$) combinatorially equivalent to a 3-polytope?

Question 2: Which self-dual 3-polytopes are combinatorially equivalent to $\mathcal{B}(V)$ for some extremal V ?

Here we insert a third question related to Question 1.

Question 3: For which extremal sets $V \subset \mathbb{R}^3$ is $\mathcal{B}(V)$ 3-polytopal, i.e., has a face structure combinatorially equivalent to that of a 3-polytope?

The answer to the first question is “no” merely because $\mathcal{B}(V)$ may have a digonal facet F_x for each dangling vertex x . So let us modify the question, by assuming that $\mathcal{B}(V)$ has no dangling vertices.

Question 1’: Is every Ball polytope $\mathcal{B}(V)$ of a *critical* (extremal) set $V \subset \mathbb{R}^3$ ($4 \leq \#V < \infty$) (i.e., V is extremal having no dangling vertices; cf. Proposition – Definition 2.1 above) combinatorially equivalent to an ordinary 3-polytope?

The answer to this modified question is again “no”: as mentioned in § 1 above, there is a critical configuration $V \subset \mathbb{R}^3$, $\#V = 8$, for which $\mathcal{B}(V)$ is not polytopal (in fact, $\text{skel}_1(\mathcal{SF}(\mathcal{B}(V)))$ is not even 3-connected). This configuration will be described in the announced subsequent paper.

An obvious partial answer to Question 2 is: a necessary condition for a 3-polytope P to be combinatorially equivalent to $\mathcal{B}(V)$ for some extremal set $V \subset \mathbb{R}^3$ ($4 \leq \#V < \infty$) is that P is *SSD*. We conjecture that this goes vice versa, formulating

Conjecture 9.1: A (necessary and) sufficient condition for the situation that the face structure $\mathcal{F}(P)$ of a 3-polytope P be isomorphic to $\mathcal{SF}(\mathcal{B}(V))$ for some extremal (in fact: critical) configuration $V \subset \mathbb{R}^3$ is that P is *SSD*.

The following conjecture answers Question 3 (provided it is true).

Conjecture 9.2: An extremal set $V \subset \mathbb{R}^3$ has a polytopal Ball polytope $\mathcal{B}(V)$ (i.e., $\mathcal{B}(V)$ is combinatorially equivalent to an ordinary 3-polytope) if and only if V is strongly critical, i.e., it has no proper subset which is extremal; cf. Definition 2.2 above.

Clearly, a strongly critical set has no dangling vertex, hence “strongly critical” $\Rightarrow$ “critical”. Good reasons for why to believe in the truth of Conjecture 9.2 (perhaps even a proof of it) will be given in the mentioned subsequent paper.

For the following conjecture we remind that the *polar body* P^Δ of a body $P \subset \mathbb{R}^d$ with $o \in \text{int}P$ is $P^\Delta =_{\text{def}} \{y \in \mathbb{R}^d : \langle y, x \rangle \leq 1 \text{ for } x \in P\}$, and that if P is a d -polytope with $o \in \text{int}P$, then the face complexes $\mathcal{F}(P^\Delta)$ of P^Δ and $\mathcal{F}(P)$ of P are dual.

Conjecture 9.3: Let Q be an *SSD* 3-polytope. Then there is a 3-polytope P combinatorially equivalent to Q such that $P^\Delta = -P$.

Acknowledgement: The authors wish to thank Dr. Margarita Spirova (Chemnitz) for preparing the figures.

References

- [1] J. ASHLEY, B. GRÜNBAUM, G. C. SHEPHARD, W. STROMQUIST: Self-duality groups and ranks of self-duality. In: *Applied Geometry and Discrete Mathematics – The Victor Klee Festschrift*, Eds.: P. Gritzmann and B. Sturmfels, DIMACS Ser. Discrete Math. Theoret. Comput. Sci. **4** (1991), 11-50.
- [2] K. BEZDEK, Z. LANGI, M. NASZÓDI, P. PAPEZ: Ball-polyhedra. *Discrete Comput. Geom.* **38** (2007), Special issue dedicated to the memory of László Fejes Tóth, 201-230.
- [3] K. BEZDEK, M. NASZÓDI: Rigidity of ball-polyhedra in Euclidean 3-space. *European J. Combin.* **27** (2006), 255-268.
- [4] V. BOLTYANSKI, H. MARTINI, P. SOLTAN: *Excursions into Combinatorial Geometry*. Springer, Berlin-Heidelberg, 1997.

- [5] P. BRASS: On the maximum number of unit distances among n points in dimension four. In: *Intuitive Geometry*, Eds.: I. Bárány et al., Bolyai Soc. Math. Studies **4** (1997), 277-290.
- [6] G. D. CHAKERIAN, H. GROEMER: Convex bodies of constant width. In: *Convexity and its Applications*, Eds. P. M. Gruber and J. M. Wills, Birkhäuser, Basel, 1983, 49-96.
- [7] V. L. DOL'NIKOV: Some properties of graphs of diameters. *Discrete Comput. Geom.* **24** (2000), 293-299.
- [8] H. G. EGGLESTON: *Convexity*. Cambridge University Press, 1958.
- [9] H. G. EGGLESTON: Sets of constant width in finite dimensional Banach spaces. *Israel J. Math.* **3** (1965), 163-172.
- [10] P. ERDŐS: On sets of distances of n points. *Amer. Math. Monthly* **53** (1946), 248-250.
- [11] P. ERDŐS: On sets of distances of n points in Euclidean space. *Magyar Tudom. Akad. Matem. Kut. Int. Közl.* (Publ. Math. Inst. Hung. Acad. Sci.) **5** (1960), 165-169.
- [12] P. ERDŐS, J. PACH: Variations on the theme of repeated distances. *Combinatorica* **10** (1990), 261-269.
- [13] B. GRÜNBAUM: A proof of Vázsonyi's conjecture. *Bull. Research Council Israel*, Section A 6 (1956), 77-78.
- [14] B. GRÜNBAUM: *Convex Polytopes*. Interscience Publishers, London-New York-Sydney, 1967 (Completed version: Springer, 2003).
- [15] B. GRÜNBAUM, G. C. SHEPHARD: Is self-duality involutory? *Amer. Math. Monthly* **95** (1988), 729-733.
- [16] A. HEPPE: Beweis einer Vermutung von A. Vázsonyi. *Acta Math. Acad. Sci. Hungar.* **7** (1956), 463-466.
- [17] H. W. E. JUNG: Über die kleinste Kugel, die eine räumliche Figur einschließt. *J. Reine Angew. Math.* **123** (1901), 241-257.

- [18] J. PACH, P. K. AGARWAL: *Combinatorial Geometry*. John Wiley & Sons, 1995.
- [19] Y. S. KUPITZ *Extremal Problems of Combinatorial Geometry*. Aarhus University, Lecture Notes Series **53** (1979).
- [20] Y. S. KUPITZ, H. MARTINI: On the weak circular intersection property. *Studia Sci. Math. Hungar.* **36** (2000), 371-385.
- [21] Y. S. KUPITZ, H. MARTINI, M. A. PERLES: Finite sets in $\mathbb{R}^d$ with many diameters – a survey. In: *Proceedings of the International Conference on Mathematics and Applications* (ICMA-MU 2005, Bangkok), Mahidol University Press, Bangkok, 2005, 91-112. Reprinted in a special volume of the *East-West J. Math.: Contributions in Mathematics and Applications* (2007), 41-57 (Zbl. 1141.51300).
- [22] Y. S. KUPITZ, H. MARTINI, B. WEGNER: A linear-time construction of Reuleaux polygons. *Beiträge Algebra Geom.* **37** (1996), 415-427.
- [23] A. E. MAYER: Eine Überkonvexität. *Math. Z.* **39** (1934) 512-531.
- [24] C. C. NEADERHAUSER, G. B. PURDY: On finite sets in E^k in which the diameter is frequently achieved. *Period. Math. Hungar.* **13** (1982), 253-257.
- [25] G. T. SALLEE: Reuleaux polytopes. *Mathematika* **17** (1970), 315-323.
- [26] Z. SCHUR, M. A. PERLES, H. MARTINI, Y. S. KUPITZ: On the number of maximal regular simplices determined by n points in $\mathbb{R}^d$. In: *Discrete and Computational Geometry – The Goodman-Pollack Festschrift*, Springer, 2003, 767-787.
- [27] S. STRASZEWICZ: Sur un problème géométrique de P. Erdős. *Bull. Acad. Polon. Sci. Cl. III*, **5** (1957), 39-40, IV-V.
- [28] P. VAN WAMELEN: The maximum number of unit distances among n points in dimension four. *Beiträge Algebra Geom.* **40** (1990), 475-477.

- [29] D. R. WOODALL: Thrackles and Deadlock. In: *Combinatorics, Proc. Conf. Comb. Math.* (D. Welsh, ed.), Academic Press, London 1971, 335-347.
- [30] I. M. YAGLOM, V. BOLTYANSKI: *Convex Figures* (transl. from Russian). New York, 1961.

Yaakov S. Kupitz

Micha A. Perles

Institute of Mathematics

The Hebrew University of Jerusalem

Jerusalem

ISRAEL

kupitz@math.huji.ac.il

perles@math.huji.ac.il

Horst Martini

Faculty of Mathematics

University of Technology Chemnitz

09107 Chemnitz

GERMANY

martini@mathematik.tu-chemnitz.de